

A SIMPLE PROOF OF PERELMAN'S COLLAPSING THEOREM FOR 3-MANIFOLDS

JIANGUO CAO AND JIAN GE

ABSTRACT. We will simplify earlier proofs of Perelman's collapsing theorem for 3-manifolds given by Shioya-Yamaguchi [SY00]-[SY05] and Morgan-Tian [MT08]. A version of Perelman's collapsing theorem states: "*Let $\{M_i^3\}$ be a sequence of compact Riemannian 3-manifolds with curvature bounded from below by (-1) and $\text{diam}(M_i^3) \geq c_0 > 0$. Suppose that all unit metric balls in M_i^3 have very small volume at most $v_i \rightarrow 0$ as $i \rightarrow \infty$ and suppose that either M_i^3 is closed or has possibly convex incompressible toral boundary. Then M_i^3 must be a graph-manifold for sufficiently large i .*" This result can be viewed as an extension of the implicit function theorem. Among other things, we apply Perelman's critical point theory (e.g., multiple conic singularity theory and his fibration theory) to Alexandrov spaces to construct the desired local Seifert fibration structure on collapsed 3-manifolds.

The verification of Perelman's collapsing theorem is the last step of Perelman's proof of Thurston's Geometrization Conjecture on the classification of 3-manifolds. A version of Geometrization Conjecture asserts that any closed 3-manifold admits a *piecewise locally homogeneous metric*. Our proof of Perelman's collapsing theorem is accessible to advanced graduate students and non-experts.

CONTENTS

0. Introduction	2
1. Brief introduction to Perelman's MCS theory and applications to local Seifert fibration	12
2. Exceptional orbits, Cheeger-Gromoll-Perelman soul theory and Perelman-Sharafutdinov flows	22
3. Admissible decompositions for collapsed 3-manifolds	40

In this version, we improve the exposition of our arguments in the earlier arXiv version. Among other things, we highlighted the use of *Perelman's version* of critical point theory for distance functions. The first author was supported in part by Nanjing University and an NSF grant.

4. Collapsing with spherical and toral fibers	44
5. Collapsing to boundary points or endpoints of the limiting space	47
6. Gluing local fibration structures and Cheeger-Gromov's compatibility condition	59
References	69

0. INTRODUCTION

In this paper, we present a simple proof of Perelman's collapsing theorem for 3-manifolds (cf. Theorem 7.4 of [Per03a]), which Perelman used to verify Thurston's Geometrization Conjecture on the classification of 3-manifolds.

0.1. Statement of Perelman's collapsing theorem and the main difficulties in its proof.

Theorem 0.1 (Perelman's Collapsing Theorem). *Let $\{(M_\alpha^3, g_{ij}^\alpha)\}_{\alpha \in \mathbb{Z}}$ be a sequence of compact oriented Riemannian 3-manifolds, closed or with convex incompressible toral boundary, and let $\{\omega_\alpha\}$ be a sequence of positive numbers with $\omega_\alpha \rightarrow 0$. Suppose that*

(1) *for each $x \in M_\alpha^3$ there exists a radius $\rho = \rho_\alpha(x)$, $0 < \rho < 1$, not exceeding the diameter of the manifold, such that the metric ball $B_{g^\alpha}(x, \rho)$ in the metric g_{ij}^α has volume at most $\omega_\alpha \rho^3$ and sectional curvatures of g_{ij}^α at least $-\rho^{-2}$;*

(2) *each component of toral boundary of M_α^3 has diameter at most ω_α , and has a topologically trivial collar of length one and the sectional curvatures of M_α^3 are between $(-\frac{1}{4} - \varepsilon)$ and $(-\frac{1}{4} + \varepsilon)$.*

Then, for sufficiently large α , M_α^3 is diffeomorphic to a graph-manifold.

If a 3-manifold M^3 admits an almost free circle action, then we say that M^3 admits a *Seifert fibration* structure or *Seifert fibred*. A *graph-manifold* is a compact 3-manifold that is a connected sum of manifolds each of which is either diffeomorphic to the solid torus or can be cut apart along a finite collection of incompressible tori into Seifert fibred 3-manifolds.

Perelman indeed listed an extra assumption (3) in Theorem 0.1 above. However, Perelman (cf. page 20 of [Per03a]) also pointed out that, if the proof of his stability theorem (cf. [Kap07]) is available, then his extra smoothness assumption (3) is in fact redundant.

The conclusion of Theorem 0.1 fails if the assumption of toral boundary is removed. For instance, the product 3-manifold $S^2 \times [0, 1]$ of a

2-sphere and an interval can be collapsed while keeping curvatures non-negative. However, $M^3 = S^2 \times [0, 1]$ is not a graph-manifold.

Under an assumption on both upper and lower bound of curvatures $-\frac{C}{\rho^2} \leq \text{curv}_{M_\alpha^3} \leq \frac{C}{\rho^2}$, the collapsed manifold M_α^3 above admits an F -structure of positive rank, by the Cheeger-Gromov collapsing theory (cf. [CG90]). It is well-known that a 3-manifold M_α^3 admits an F -structure of positive rank if and only if M_α^3 is a graph-manifold, (cf. [Rong93]).

On page 153 of [SY05], Shioya and Yamaguchi stated a version of Theorem 0.1 above for the case of closed manifolds, but their proof works for case of manifolds with incompressible convex toral boundary as well. Morgan and Tian [MT08] presented a proof of Theorem 0.1 without assumptions on diameters but with Perelman's extra smoothness assumption (3) which we discuss below.

It turned out that the diameter assumption is related to the study of diameter-collapsed 3-manifolds with curvature bounded from below. To see this relation, we state a local re-scaled version of Perelman's collapsing Theorem 0.1.

Theorem 0.1'. (Re-scaled version of Theorem 0.1) *Let $\{(M_\alpha^3, g_{ij}^\alpha)\}_{\alpha \in \mathbb{Z}}$ be and let $\{\omega_\alpha\}$ be a sequence of positive numbers as in Theorem 0.1 above, $x_\alpha \in M_\alpha^3$ and $\rho_\alpha = \rho_\alpha(x_\alpha)$. Suppose that there exists a re-scaled family of pointed Riemannian manifolds $\{(M_\alpha^3, \rho_\alpha^{-2} g_{ij}^\alpha), x_\alpha\}_{\alpha \in \mathbb{Z}}$ satisfying the following conditions:*

- (i) *The re-scaled Riemannian manifold $(M_\alpha^3, \rho_\alpha^{-2} g_{ij}^\alpha)$ has $\text{curv} \geq -1$ on the ball $B_{\rho_\alpha^{-2} g_{ij}^\alpha}(x_\alpha, 1)$;*
- (ii) *The diameters of the re-scaled manifolds $\{(M_\alpha^3, \rho_\alpha^{-2} g_{ij}^\alpha)\}$ are uniformly bounded from below by 1; i.e.;*

$$\text{diam}(M_\alpha^3, \rho_\alpha^{-2} g_{ij}^\alpha) \geq 1 \tag{0.1}$$

- (iii) *The volumes of unit metric balls collapse to zero, i.e.:*

$$\text{Vol}[B_{\rho_\alpha^{-2} g_{ij}^\alpha}(x_\alpha, 1)] \leq \omega_\alpha \rightarrow 0,$$

as $\alpha \rightarrow \infty$.

Then, for sufficiently large α , the collapsing 3-manifold M_α^3 is a graph-manifold.

Without inequality (0.1) above, the volume-collapsing 3-manifolds could be diameter-collapsing. Perelman's condition (0.1) ensures that the normalized family $\{(M_\alpha^3, \rho_\alpha^{-2} g_{ij}^\alpha)\}$ can *not* collapse to a point uniformly. By *collapsing to a point uniformly*, we mean that there is an additional family of scaling constants $\{\lambda_\alpha\}$ such that the sequence

$\{(M_\alpha^3, \lambda_\alpha g_{ij}^\alpha)\}$ is convergent to a 3-dimensional (possibly singular) manifold Y_∞^3 with non-negative curvature. Professor Karsten Grove kindly pointed out that the study of diameter-collapsing theory for 3-manifolds might be related to a weak version of the Poincaré conjecture. For this reason, Shioya and Yamaguchi made the following conjecture.

Conjecture 0.2 (Shioya-Yamaguchi [SY00] page 4). *Suppose that Y_∞^3 is a 3-dimensional compact, simply-connected, non-negatively curved Alexandrov space without boundary and that Y_∞^3 is a topological manifold. Then Y_∞^3 is homeomorphic to a sphere.*

Shioya and Yamaguchi ([SY00], page 4) commented that *if Conjecture 0.2 is true then the study of collapsed 3-manifolds with curvature bounded from below would be completely understood*. They also observed that Conjecture 0.2 is true for a special case when the closed (possibly singular) manifold Y_∞^3 above is a *smooth* Riemannian manifold with non-negative curvature; this is due to Hamilton's work on 3-manifolds with non-negative Ricci curvature (cf. [Ham86]).

Coincidentally, Perelman added the extra smoothness assumption (3) in his collapsing theorem.

Perelman's Smoothness Assumption 0.3 (cf. [Per03a]). *For every $w' > 0$ there exist $r = r(w') > 0$ and constants $K_m = K_m(w') < \infty$ for $m = 0, 1, 2, \dots$, such that for all α sufficiently large, and any $0 < r \leq \bar{r}$, if the ball $B_{g_\alpha}(x, r)$ has volume at least $w'r^3$ and sectional curvatures at least $-r^{-2}$, then the curvature and its m -th order covariant derivatives at x are bounded by $K_0 r^{-2}$ and $K_m r^{-m-2}$ for $m = 1, 2, \dots$, respectively.*

Let us explain how Perelman's Smoothness Assumption 0.3 is related to the smooth case of Conjecture 0.2 and let $\{((M_\alpha^3, \rho_\alpha^{-2} g_{ij}^\alpha), x_\alpha)\}_{\alpha \in \mathbb{Z}}$ be as in Theorem 0.1'. If we choose the new scaling factor λ_α^2 such that $\lambda_\alpha^2 / \rho_\alpha^{-2} \rightarrow +\infty$ as $\alpha \rightarrow \infty$, then the newly re-scaled metric $\lambda_\alpha^2 g_{ij}^\alpha$ will have sectional curvature $\geq -\frac{\rho_\alpha^{-2}}{\lambda_\alpha^2} \rightarrow 0$ as $\alpha \rightarrow \infty$. Suppose that (Y_∞, y_∞) is a pointed Gromov-Hausdorff limit of a subsequence of $\{((M_\alpha^3, \lambda_\alpha^2 g_{ij}^\alpha), x_\alpha)\}_{\alpha \in \mathbb{Z}}$. Then the limiting metric space Y_∞ will have non-negative curvature and a possibly singular metric. When $\dim[Y_\infty] = 3$, by Perelman's Smoothness Assumption 0.3, the limiting metric space Y_∞ is indeed a *smooth* Riemannian manifold of non-negative curvature. In this smooth case, Conjecture 0.2 is known to be true, (see [Ham86]).

For simplicity, we let $\hat{g}_{ij}^\alpha = \rho_\alpha^{-2} g_{ij}^\alpha$ be as in Theorem 0.1'. By Gromov's compactness theorem, there is a subsequence of a pointed Riemannian manifolds $\{((M_\alpha^3, \hat{g}_{ij}^\alpha), x_\alpha)\}$ convergent to a lower dimensional

pointed space (X^k, x_∞) of dimension either 1 or 2, i.e.:

$$1 \leq \dim[X^k] \leq 2 \quad (0.2)$$

using (0.1). To establish Theorem 0.1, it is important to establish that, for sufficiently large α , the collapsed manifold M_α^3 has a decomposition $M_\alpha^3 = \cup_i U_{\alpha,i}$ such that each $U_{\alpha,i}$ admits an almost-free circle action:

$$S^1 \rightarrow U_{\alpha,i} \rightarrow X_{\alpha,i}^2$$

We also need to show that these almost-free circle actions are compatible (almost commute) on possible overlaps.

Let us first recall how Perelman's collapsing theorem for 3-manifolds plays an important role in his solution to Thurston's Geometrization Conjecture on the classification of 3-manifolds.

0.2. Applications of collapsing theory to the classification of 3-manifolds.

In 2002-2003, Perelman posted online three important but densely written preprints on Ricci flows with surgery on compact 3-manifolds ([Per02], [Per03a] and [Per03b]), in order to solve both the Poincaré conjecture and Thurston's conjecture on Geometrization of 3-dimensional manifolds. Thurston's Geometrization Conjecture states that *"for any closed, oriented and connected 3-manifold M^3 , there is a decomposition $[M^3 - \bigcup \Sigma_j^2] = N_1^3 \cup N_2^3 \cdots \cup N_m^3$ such that each N_i^3 admits a locally homogeneous metric with possible incompressible boundaries Σ_j^2 , where Σ_j^2 is homeomorphic to a quotient of a 2-sphere or a 2-torus"*. There are exactly 8 homogeneous spaces in dimension 3. The list of 3-dimensional homogeneous spaces includes 8 geometries: $\mathbb{R}^3$, $\mathbb{H}^3$, $\mathbf{S}^3$, $\mathbb{H}^2 \times \mathbb{R}$, $\mathbf{S}^2 \times \mathbb{R}$, $\tilde{SL}(2, \mathbb{R})$, *Nil* and *Sol*.

Thurston's Geometrization Conjecture suggests the existence of especially nice metrics on 3-manifolds and consequently, a more analytic approach to the problem of classifying 3-manifolds. Richard Hamilton formalized one such approach by introducing the Ricci flow equation on the space of Riemannian metrics:

$$\frac{\partial g(t)}{\partial t} = -2\text{Ric}(g(t)) \quad (0.3)$$

where $\text{Ric}(g(t))$ is the Ricci curvature tensor of the metric $g(t)$. Beginning with any Riemannian manifold (M, g_0) , there is a solution $g(t)$ of this Ricci flow on M for t in some interval such that $g(0) = g_0$. In dimension 3, the fixed points (up to re-scaling) of this equation include the Riemannian metrics of constant Ricci curvature. For instance, they are quotients of $\mathbb{R}^3$, $\mathbb{H}^3$ and $\mathbf{S}^3$ up to scaling factors. It is easy to see

that, on compact quotients of $\mathbb{R}^3$ or $\mathbb{H}^3$, the solution to Ricci flow equation is either stable or expanding. Thus, on compact quotients of $\mathbb{R}^3$ and $\mathbb{H}^3$, the solution to Ricci flow equation exists for all time $t \geq 0$.

However, on quotients of $\mathbf{S}^2 \times \mathbb{R}$ or $\mathbf{S}^3$, the solution $\{g(t)\}$ to Ricci flow equation exists only for finite time $t < T_0$. Hence, one knows that in general the Ricci flow will develop singularities in finite time, and thus a method for analyzing these singularities and continuing the flow past them must be found.

These singularities may occur along proper subsets of the manifold, not the entire manifold. Thus, Perelman introduced a more general evolution process called Ricci flow with surgery (cf. [Per02] and [Per03a]). In fact, a similar process was first introduced by Hamilton in the context of four-manifolds. This evolution process is still parameterized by an interval in time, so that for each t in the interval of definition there is a compact Riemannian 3-manifold M_t . However, there is a discrete set of times at which the manifolds and metrics undergo topological and metric discontinuities (surgeries). Perelman did surgery along 2-spheres rather than surfaces of higher genus, so that the change in topology for $\{M_t\}$ turns out to be completely understood. More precisely, Perelman's surgery on 3-manifolds is *the reverse process of taking connected sums*: cut a 3-manifold along a 2-sphere and then cap-off by two disjoint 3-balls. Perelman's surgery processes produced exactly the topological operations needed to cut the manifold into pieces on which the Ricci flow can produce the metrics sufficiently controlled so that the topology can be recognized. It was expected that each connected components of resulting new manifold M_t^3 is either a graph-manifold or a quotient of one of homogeneous spaces listed above. It is well-known that any graph-manifold is a union of quotients of 7 (out of the 8 possible) homogeneous spaces described above. More precisely, Perelman presented a very densely written proof of the following result.

Theorem 0.4 (Perelman [Per02],[Per03a], [Per03b]). *Let (M^3, g_0) be a closed and oriented Riemannian 3-manifold. Then there is a Ricci flow with surgery, say $\{(M_t, g(t))\}$, defined for all $t \in [0, T)$ with initial metric (M, g_0) . The set of discontinuity times for this Ricci flow with surgery is a discrete subset of $[0, \infty)$. The topological change in the 3-manifold as one crosses a surgery time is a connected sum decomposition together with removal of connected components, each of which is diffeomorphic to one of $(S^2 \times S^1)/\Gamma_i$ or S^3/Γ_j . Furthermore, there are two possibilities:*

(1) *Either the Ricci flow with surgery terminates at a finite time T . In this case, M_0^3 is diffeomorphic to the connect sum of $(S^2 \times$*

$S^1)/\Gamma_i$ and S^3/Γ_j . In particular, if M_0^3 is simply-connected, then M_0^3 is diffeomorphic to S^3 .

(2) Or the Ricci flow with surgery exists for all time, i.e., $T = \infty$.

The detailed proof of Perelman's theorem above can be found in [CaZ06], [KL08] and [MT07]. In fact, if M^3 is a simply-connected closed manifold, then using a theorem of Hurwicz, one can find that $\pi_3(M^3) = H_3(M^3, \mathbb{Z}) = \mathbb{Z}$. Hence, there is a map $F : S^3 \rightarrow M^3$ of degree 1. One can view S^3 as a two-point suspension of S^2 , i.e., $S^3 \sim S^2 \times [0, 1]/\{0, 1\}$. Thus, for such a manifold M^3 with a metric g , one can define the 2-dimensional width $W_2(M, g) = \inf_{\deg(F)=1} \max_{0 \leq s \leq 1} \{Area_g[F(S^2, s)]\}$, (compare with [CalC92]). Colding and Minicozzi (cf. [CM05], [CM08a], [CM08b]) established that

$$0 < W_2(M_t^3, g(t)) \leq (t + c_1)^{\frac{3}{4}} [c_2 - 16\pi(t + c_3)^{\frac{1}{4}}] \quad (0.4)$$

for Perelman's solutions $\{M_t^3, g(t)\}$ to Ricci flow with surgery, where $\{c_1, c_2, c_3\}$ are positive constants independent of $t \in [0, T)$. Therefore, for a simply-connected closed manifold M_0^3 , it follows from (0.4) that the Ricci flow with surgery must end at a finite time T . Thus, by Theorem 0.4, the conclusion of the Poincaré conjecture holds for such a simply-connected closed 3-manifold M_0^3 . Other proofs of Perelman's finite time extinction theorem can be found in [Per03b] and [MT07].

It remains to discuss the long time behavior of Ricci curvature flow with surgery. We will perform the so-called Margulis thick-thin decomposition for a complete Riemannian manifold (M^n, g) . The thick part is the *non-collapsing part* of (M^n, g) , while the thin part is the *collapsing* portion of (M^n, g) .

Let $\rho(x, t)$ denote the radius ρ of the metric ball $B_{g(t)}(x, \rho)$, where we may choose $\rho(x, t)$ so that

$$\inf\{\sec_{g(t)}|_y \mid y \in B_{g(t)}(x, \rho)\} \geq -\rho^{-2}$$

and $\frac{1}{2}\rho(x, t) \leq \rho(y, t) \leq \rho(x, t)$ for all $y \in B_{g(t)}(x, \rho/2)$. The re-scaled thin part of M_t can be defined as

$$M_-(\omega, t) = \{x \in M_t^3 \mid \text{Vol}[B_{g(t)}(x, \rho(x, t))] < \omega\rho^3(x, t)\} \quad (0.5)$$

and its complement is denoted by $M_+(\omega, t)$, which is called the thick part for a fixed positive number ω .

When there is no surgery occurring for all time $t \geq 0$, Hamilton successfully classified the thick part $M_+(\omega, t)$.

Theorem 0.5 (Hamilton [Ham99], non-collapsing part). *Let $\{(M_t^3, g(t))\}$, $M_-(\omega, t)$ and $M_+(\omega, t)$ be as above. Suppose that M_t^3 is diffeomorphic to M_0^3 for all $t \geq 0$. Then there are only two possibilities:*

- (1) If there is no thin part (i.e., $M_-(\omega, t) = \emptyset$), then either $(M_t^3, g(t))$ is convergent to a flat 3-manifold or $(M_t^3, \frac{2}{t}g(t))$ is convergent to a compact quotient of hyperbolic space $\mathbb{H}^3$;
- (2) If both $M_+(\omega, t)$ and $M_-(\omega, t)$ are non-empty, then the thick part $M_+(\omega, t)$ is diffeomorphic to a disjoint union of quotients of real hyperbolic space $\mathbb{H}^3$ with finite volume and with cuspidal ends removed.




FIGURE 1. 2-dimensional thick-thin decomposition




FIGURE 2. 3-dimensional thick-thin decomposition

Perelman (cf. [Per03a]) asserted that the conclusion of Theorem 0.5 holds if we replace the classical Ricci flow by *the Ricci flow with surgeries*, (cf. [Per03a]). Detailed proof of this assertion of Perelman can be found in [CaZ06], [KL08] and [MT10].

Suppose that $\mathbb{H}^3/\Gamma$ is a complete but non-compact hyperbolic 3-manifold with finite volume. The cuspidal ends of $\mathbb{H}^3/\Gamma$ are exactly the thin parts of $M_\infty^3 = \mathbb{H}^3/\Gamma$. Each cuspidal end of $\mathbb{H}^3/\Gamma$ is diffeomorphic to a product of a torus and half-line (i.e., $T^2 \times [0, \infty)$). Hence, each cusp is a graph-manifold.

It should be pointed out that possibly infinitely many surgeries took place *only* on thick parts of manifolds $\{M_t\}$ after appropriate re-scalings, due to the celebrated Perelman's κ -non-collapsing theory, (see [Per02]).

Moreover, Perelman (cf. [Per03a]) pointed out that the study of the thin part $M_-(\omega, t)$ has nothing to do with Ricci flow, but is related

to *his version* of critical point theory for distance functions. We now outline our simple proof of Theorem 0.1 using Perelman's version of critical point theory in next sub-section.

0.3. Outline of a proof of Perelman's collapsing theorem.

In order to illustrate main strategy in the proof of Perelman's collapsing theorem for 3-manifolds, we make some general remarks. Roughly speaking, Perelman's collapsing theorem can be viewed as a generalization of the implicit function theorem. Suppose that $\{M_\alpha^3\}$ is a sequence of collapsing 3-manifolds with curvature ≥ -1 and that $\{M_\alpha^3\}$ is not a diameter-collapsing sequence. To verify that M_α^3 is a graph-manifold for sufficiently large α , it is sufficient to construct a decomposition $M_\alpha^3 = \cup_{i=1}^{m_\alpha} U_{\alpha,i}$ and a collection of *regular* functions (or maps) $F_{\alpha,i} : U_{\alpha,i} \rightarrow \mathbb{R}^{s_i}$, where $s_i = 1$ or 2 . We require that the collection of locally defined functions (or maps) $\{(U_{\alpha,i}, F_{\alpha,i})\}_{i=1}^{m_\alpha}$ satisfy two conditions:

- (i) Each function (or map) $F_{\alpha,i}$ is *regular* enough so that Perelman's version of implicit function theorem (cf. Theorem 1.2 below) is applicable;
- (ii) The collection of locally defined *regular* functions (or maps) are compatible on any possible overlaps in the sense of Cheeger-Gromov (cf. [CG86] [CG90]). More precisely, if $U_{\alpha,i} \cap U_{\alpha,j} \neq \emptyset$ and if $[F_{\alpha,i}^{-1}(y) \cap F_{\alpha,j}^{-1}(z)] \neq \emptyset$ with $\dim[F_{\alpha,i}^{-1}(y)] \leq \dim[F_{\alpha,j}^{-1}(z)]$, then we require that either $F_{\alpha,i}^{-1}(y) \subset F_{\alpha,j}^{-1}(z)$ or the union $[F_{\alpha,i}^{-1}(y) \cup F_{\alpha,j}^{-1}(z)]$ is contained in a 2-dimensional orbit of an almost-free torus action.

If the above two conditions are met, with additional efforts, we can construct a *compatible* family of the locally defined Seifert fibration structures (which is equivalent to an F -structure $\mathcal{F}$ of positive rank in the sense of Cheeger-Gromov) on a sufficiently collapsed 3-manifold M_α^3 . It follows that M_α^3 is a graph-manifold for sufficiently large α , (cf. [Rong93]).

Perelman's choices of locally defined *regular* functions (or maps) are related to distance functions $r_{A_{\alpha,i}}(x) = d(x, A_{\alpha,i})$ from appropriate subsets $A_{\alpha,i}$. We briefly illustrate the main strategy of our proof for the following two cases.

Case 1. *The metric balls $\{B_{M_\alpha^3}(x_\alpha, r)\}$ collapse to an open interval.*

We will show that $B_{M_\alpha^3}(x_\alpha, r)$ is homeomorphic to a slim cylinder $N_\alpha^2 \times I$ with shrinking spherical or toral factor N_α^2 for sufficiently large α . When a sequence of the pointed 3-manifolds $\{(B_{M_\alpha^3}(x_\alpha, r), x_\alpha)\}$ with curvature ≥ -1 are convergent to an 1-dimensional space (X^1, x_∞)

and x_∞ is an interior point, Perelman-Yamaguchi fibration theory is applicable. Thus, we are led to consider the fibration

$$N_\alpha^2 \rightarrow B_{M_\alpha^3}(x_\alpha, r) \xrightarrow{F_\alpha} (-\ell, \ell)$$




FIGURE 3. Slim cylinders with collapsing fibers.

We now discuss the topological type of the fiber N_α^2 . Let $x_\infty \in (-\ell, \ell)$ and ε_α be the diameter of $F_\alpha^{-1}(x_\infty)$ in M_α^3 . We further consider the limiting space Y_∞^s of re-scaled spaces $\{(B_{\varepsilon_\alpha^{-1}M_\alpha^3}(x_\alpha, \frac{r}{\varepsilon_\alpha}), x_\alpha)\}$, as $\varepsilon_\alpha \rightarrow 0$. There are two sub-cases: $\dim(Y_\infty^s) = 3$ or $\dim(Y_\infty^s) = 2$. Let us consider the subcase of $\dim(Y_\infty^s) = 3$:


$$N_\infty^2 \rightarrow Y_\infty^3 \rightarrow \mathbb{R}$$

where both N_∞^2 and Y_∞^3 are manifolds with possibly singular metrics of non-negative curvature. To classify singular surfaces N_∞^2 with non-negative curvature, we use a splitting theorem and the distance non-increasing property of Perelman-Sharafutdinov retraction on the universal cover $\tilde{N}_\infty^2$, when N_∞^2 has non-zero genus. With some extra efforts, we will conclude that N_∞^2 must be homeomorphic to a quotient of 2-sphere or 2-torus, (see Section 2 below). It now follows from a version of Perelman's stability theorem that the fiber N_α^2 is homeomorphic to N_∞^2 , for sufficiently large α . Hence, N_α^2 is a quotient of 2-sphere or 2-torus as well, for sufficiently large α . Our new proof of Perelman's collapsing theorem for this subcase is much simpler than the approach of Shioya-Yamaguchi presented in [SY00].

The sub-case of $\dim(Y_\infty^s) = 2$ is related to the following case:

Case 2. *The metric balls $\{B_{M_\alpha^3}(x_\alpha, r)\}$ collapse to an open disk.*

We will show that $B_{M_\alpha^3}(x_\alpha, r)$ is homeomorphic to a fat solid torus $D^2 \times S_{\varepsilon_\alpha}^1$ with shrinking core $S_{\varepsilon_\alpha}^1$ for sufficiently large α .

FIGURE 4. Fat solid tori with shrinking cores S_ϵ^1 .

In this case, our strategy can be illustrated in the following diagram

$$\begin{array}{ccc}
 B_{M_\alpha^3}(x_\alpha, \delta) & \xrightarrow{F_\alpha} & \mathbb{R}^2 \\
 \downarrow \text{G-H} & & \uparrow \\
 B_{X^2}(x_\infty, \delta) & \xrightarrow{F_\infty} & \mathbb{R}^2
 \end{array}$$

where the sequence of metric balls $B_{(M_\alpha^3, \hat{g}^\alpha)}(x_\alpha, \delta)$ are convergent to the metric disk $B_{X^2}(x_\infty, \delta)$ for $\delta \leq r$. We will construct an admissible map F_∞ which is regular at the punctured disk, using Perelman's multiple conic singularity theory, (see Theorem 1.17 below). Among other things, we will use the following result of Perelman to construct the desired map F_∞ .

Theorem 0.6 (Conic Lemma in Perelman's critical point theory, (cf. [Per94] page 211)). *Let X be an Alexandrov space of dimension k , curv ≥ -1 and $x \in X$ be an interior point of X . Then the distance function $r_x(y) = d(x, y)$ has no critical points on $[B_X(x, \delta) - \{x\}]$ for sufficiently small δ depending on x .*

We will use Theorem 0.6 and Perelman's semi-flow orbit stability theorem (cf. Proposition 1.14 below) to conclude that the lifting maps F_α is regular on the annular region $A_{M_\alpha^3}(x_\alpha, \epsilon, \delta) = [B_{M_\alpha^3}(x_\alpha, \delta) - B_{M_\alpha^3}(x_\alpha, \epsilon)]$. With extra efforts, one can construct a local Seifert fibration structure:

$$S^1 \rightarrow A_{M_\alpha^3}(x_\alpha, \epsilon, \delta) \xrightarrow{G_\alpha} A_{X^2}(x_\infty, \epsilon, \delta)$$

In summary, Perelman's collapsing theorem for 3-manifolds can be viewed an extension of the implicit function theorem. Our proof of Perelman's collapsing theorem benefited from *his version* of critical point theory for distance functions, including his conic singularity theory and fibration theory. Perelman's multiple conic singularity theory

and his fibration theory are designed for possibly singular Alexandrov spaces X^k . Therefore, the smoothness of metrics on X^k does *not* play a big role in the applications of Perelman's critical point theory, unless we run into the so-called essential singularities (or extremal subsets). When essential singularities do occur on surfaces, we use the MCS theory (e.g. Theorem 0.6) and the multiple-step Perelman-Sharafutdinov flows to handle them, (see §5.2 below).

Without using Perelman's version of critical point theory, Shioya-Yamaguchi's proof of the collapsing theorem for 3-manifolds was lengthy and involved. For instance, they use their singular version of Gauss-Bonnet theorem to classify surfaces of non-negative curvature, (see Chapter 14 of [SY00]). The proof of the singular version of the Gauss-Bonnet theorem was non-trivial. In addition, Shioya-Yamaguchi extended the Cheeger-Gromoll soul theory to 3-dimensional singular spaces with non-negative curvature, which was rather technical and occupied the half of their first paper [SY00]. Using Perelman's version of critical point theory, we will provide alternative approaches to classify non-negatively curved surfaces and open 3-manifolds with possibly singular metrics, (e.g., the 3-dimensional soul theory). Our arguments inspired by Perelman are considerably shorter than Shioya-Yamaguchi's proof for the 3-dimensional soul theory, (see §2.2 below).

For the readers who prefer a traditional proof of the collapsing theorem without using Perelman's version of critical point theory, we recommend the important papers of Morgan-Tian [MT08] and Shioya-Yamaguchi [SY00], [SY05]. Finally, we should also mention the recent related work of Gerard Besson et al, (cf. [BBBMP10]). Another proof of Perelman's collapsing theorem for 3-manifolds has been announced by Kleiner and Lott (cf. [KL10]).

We refer the organization of this paper to the table of contents at the beginning. In Section 1-4 below, we mostly discuss interior points of Alexandrov spaces, unless otherwise specified.

1. BRIEF INTRODUCTION TO PERELMAN'S MCS THEORY AND APPLICATIONS TO LOCAL SEIFERT FIBRATION

In §1-2, we will discuss our proof of Theorem 0.1' for a special case. In this special case, we assume that the sequence of metric balls $\{B_{(M_\alpha^3, g^\alpha)}(x_\alpha, r)\}$ is convergent to a metric ball $B_{X^2}(x_\infty, r)$, where x_∞ is an interior point of X^2 . Using several known results of Perelman, we will show that there is a (possibly singular) circle fibration:

$$S^1 \rightarrow B_{(M_\alpha^3, g^\alpha)}(x_\alpha, \varepsilon) \rightarrow B_{X^2}(x_\infty, \varepsilon), \quad (1.1)$$

for some $\varepsilon < r$. In other words, we shall show that $\hat{B}_{M_\alpha^3}(x_\alpha, \varepsilon)$ looks like a *fat* solid tori with a shrinking core, i.e., $\hat{B}_{M_\alpha^3}(x_\alpha, \varepsilon) \sim [D^2 \times S_\varepsilon^1] \sim [(D^2 \times (\mathbb{R}/\varepsilon\mathbb{Z}))]$, (see Figure 4 above and Example 2.0 below).

In fact, using the Conic Lemma (Theorem 0.6 above), Kapovitch [Kap05] already established a circle-fibration structure over the annular region $A_{X^2}(x_\infty, \delta, \varepsilon)$. Let $\Sigma_x(X)$ denote the space of unit directions of an Alexandrov space X of curvature ≥ -1 at point x . When $\dim(X) = 2$, it is known (cf. [BGP92]) that X^2 must be a 2-dimensional topological manifold. Thus, $\Sigma_x(X^2)$ is a circle, and hence an 1-dimensional manifold.

Proposition 1.1 ([Kap05], page 533). *Suppose that $M_\alpha^n \xrightarrow{G-H} X$, where M_α^n is a sequence of n -dimensional Riemannian manifolds with sectional curvature ≥ -1 . Suppose that there exists $x_\infty \in X$ such that $\Sigma = \Sigma_{x_\infty}(X)$ is a closed Riemannian manifold. Then there exists $r_0 = r_0(x_\infty)$ such that for any $M_\alpha \ni x_\alpha \rightarrow x_\infty$ we have: For any sufficiently large α , and $r \leq r_0$, there exists a topological fiber bundle*

$$S_\alpha \rightarrow \partial B_{M_\alpha}(x_\alpha, r) \rightarrow \Sigma_{x_\infty}(X)$$

such that

- (1) S_α and $\partial B_{M_\alpha}(x_\alpha, r)$ are topological manifolds;
- (2) Both S_α and $\partial B_{M_\alpha}(x_\alpha, r)$ are connected;
- (3) The fundamental group $\pi_1(S_\alpha)$ of the fiber is almost nilpotent.

We will use Perelman's fibration theorem and an multiple conic singularity theory to establish the desired circle fibration over the annular region $A_{X^2}(x_\infty, \delta, \varepsilon)$ for $\delta < \varepsilon$. Our strategy can be illustrated in the following diagram

$$\begin{array}{ccc} B_{M_\alpha^3}(x_\alpha, r) & \xrightarrow{F_\alpha} & \mathbb{R}^2 \\ \downarrow \text{G-H} & & \downarrow \\ B_{X^2}(x_\infty, r) & \xrightarrow{F_\infty} & \mathbb{R}^2 \end{array}$$

where the sequence of metric balls $B_{(M_\alpha^3, \hat{g}^\alpha)}(x_\alpha, r)$ are convergent to the metric disk $B_{X^2}(x_\infty, r)$.

If F_α were a “topological submersion” to its image, then we would be able to obtain the desired topological fibration. For this purpose, we will recall Perelman's Fibration Theorem for non-smooth maps.

1.1. Brief introduction to Perelman's critical point theory.

We postpone the definition of admissible maps to §1.2. In §1.2, we will also recall the notion of *regular points* for a sufficiently wide class of “*admissible mappings*” from an Alexandrov space X^n to Euclidean space $\mathbb{R}^k$.

Let $F : X^n \rightarrow \mathbb{R}^k$ be an admissible map. The points of an Alexandrov space X that are not regular are said to be critical points, and their images in $\mathbb{R}^k$ are said to be critical values of $F : X \rightarrow \mathbb{R}^k$. All other points of $\mathbb{R}^k$ are called regular values.

Theorem 1.2 (Perelman's Fibration Theorem [Per94] page 207).

- (A) *An admissible mapping is open and admits a trivialization in a neighborhood of each of its regular points.*
- (B) *If an admissible mapping has no critical points and is proper in some domain, then its restriction to this domain is the projection of a locally trivial fiber bundle.*

There are several equivalent definitions of Alexandrov spaces of $\text{curv} \geq k$. Roughly speaking, a length space X is said to have curvature ≥ 0 if and only if, for any geodesic triangle Δ in X , the corresponding triangle $\tilde{\Delta}$ of the same side-lengths in $\mathbb{R}^2$ is thinner than Δ .

More precisely, let M_k^2 be a simply connected complete surface of constant sectional curvature k . A triangle in a length space X consists of three vertices, say $\{a, b, c\}$ and three length-minimizing geodesic segments $\{\overline{ab}, \overline{bc}, \overline{ac}\}$. Let $|ab|$ be the length of $\overline{ab}$. Given a real number k , a comparison triangle $\Delta_{\tilde{a}, \tilde{b}, \tilde{c}}^k$ is a triangle in M_k^2 with the same side lengths. Its angles are called the comparison angles and denoted by $\tilde{\angle}_a^k(b, c)$, etc. A comparison triangle exists and is unique whenever $k \leq 0$ or $k > 0$ and $|ab| + |bc| + |ca| < \frac{2\pi}{\sqrt{k}}$.

Definition 1.3. *A length space X is called an Alexandrov space of curvature $\geq k$ if any $x \in X$ has a neighborhood U_x for any $\{a, b, c, d\} \in U_x$, the following inequality*

$$\tilde{\angle}_a^k(b, c) + \tilde{\angle}_a^k(c, d) + \tilde{\angle}_a^k(d, b) \leq 2\pi.$$

Alexandrov spaces with $\text{curv} \geq \lambda$ have several nice properties, (cf. [BGP92]). For instance, the dimension of an Alexandrov space X is either an integer or infinite. Moreover, for any $x \in X$, there is a well defined tangent cone $T_x^-(X)$ along with an “inner product” on $T_x^-(X)$.

In fact, if X is an Alexandrov space with the metric d , then we denote by λX the space $(X, \lambda d)$. Let $i_\lambda : \lambda X \rightarrow X$ be the canonical map. The Gromov-Hausdorff limit of pointed spaces $\{(\lambda X, x)\}$ for $\lambda \rightarrow \infty$ is the tangent cone $T_x^-(X)$ at x , (see §7.8.1 of [BGP92]).

For any function $f : X \rightarrow \mathbb{R}$, the function $d_x f : T_x^-(X) \rightarrow \mathbb{R}$ such that

$$d_x f = \lim_{\lambda \rightarrow +\infty} \frac{f \circ i_\lambda - f(x)}{1/\lambda}$$

is called the differential of f at x .

Let us now recall the notion of regular points for distance functions.

Definition 1.4 ([GS77], [Gro81]). *Let $A \subset X$ be a closed subset of an Alexandrov space X and $f_A(x) = d(x, A)$ be the corresponding distance function from A . A point $x \notin A$ is said to be a regular point of f_A if there exists a non-zero direction $\vec{\xi} \in T_x^-(X)$ such that*

$$d_x f(\vec{\xi}) > 0. \quad (1.2)$$

It is well-known that if X has $\text{curv} \geq 0$ then $f(x) = \frac{1}{2}[d(x, p)]^2$ has the property that $\text{Hess}(f) \leq I$, (see [Petr07]). To explain such an inequality, we recall the notion of semi-concave functions.

Definition 1.5 ([Per94] page 210). *A function $f : X \rightarrow \mathbb{R}$ is said to be λ -concave in an open domain U if for any length-minimizing geodesic segment $\sigma : [a, b] \rightarrow U$ of unit speed, the function*

$$f \circ \sigma(t) - \lambda t^2/2$$

is concave.

When f is 1-concave, we say that $\text{Hess}(f) \leq I$. It is clear that if $f : U \rightarrow \mathbb{R}$ is a semi-concave function, then

$$d_x f : T_x^-(X) \rightarrow \mathbb{R}$$

is a concave function.

In order to introduce the notion of semi-gradient vector for a semi-concave function f , we need to recall the notion of “inner product” on $T_x^-(X)$. For any pair of vectors $\vec{u}$ and $\vec{v}$ in $T_x^-(X)$, we define

$$\langle \vec{u}, \vec{v} \rangle = \frac{1}{2}(|\vec{u}|^2 + |\vec{v}|^2 - |\vec{u}\vec{v}|^2) = |\vec{u}||\vec{v}| \cos \theta$$

where θ is the angle between $\vec{u}$ and $\vec{v}$, $|\vec{u}\vec{v}| = d_{T_x^-(X)}(\vec{u}, \vec{v})$, $|\vec{u}| = d_{T_x^-(X)}(\vec{u}, o)$ and o denotes the origin of the tangent cone.

Definition 1.6 ([Petr07]). *For any given semi-concave function f on X , a vector $\vec{\eta} \in T_x^-(X)$ is called a gradient of f at x (in short $\vec{\eta} = \nabla f$) if*

- (i) $d_x f(\vec{v}) \leq \langle \vec{\eta}, \vec{v} \rangle$ for any $\vec{v} \in T_x^-(X)$;
- (ii) $d_x f(\vec{\eta}) = |\vec{\eta}|^2$.

It is easy to see that any semi-concave function has a uniquely defined gradient vector field. Moreover, if $d_x f(\vec{v}) \leq 0$ for all $\vec{v} \in T_x^-(X)$, then $\nabla f|_x = 0$. In this case, x is called a critical point of f . Otherwise, we set

$$\nabla f = d_x f(\vec{\xi})\vec{\xi}$$

where $\vec{\xi}$ is the (necessarily unique) unit vector for which $d_x f$ attains its positive maximum on $\Sigma_x(X)$, where $\Sigma_x(X)$ is the space of direction of X at x .

Proposition 1.7 ([Per94], [PP94]). *Let X^n be a metric space with curvature ≥ -1 and $\hat{x}$ be an interior point of X^n . Then there exists a strictly concave function $h : B(\hat{x}, r) \rightarrow (-\infty, 0]$ such that (1) $h(\hat{x}) = 0$ and $B(\hat{x}, \frac{s}{\lambda}) \subset h^{-1}((-s, 0]) \subset B(\hat{x}, \lambda s)$ for $s \leq \frac{r}{4\lambda}$; (2) the distance function $r_{\hat{x}}(y)$ has no critical point in punctured ball $[B_X(\hat{x}, \varepsilon) - \{\hat{x}\}]$, for some $\{\varepsilon, r, \lambda\}$ depending on $\hat{x}$.*

Proof. (1) The construction of the strictly concave function h described above is available in literature (see [GW97], [Kap02]). In fact, let $f_{\delta'} : X \rightarrow \mathbb{R}$ be defined as on page 129 of [Kap02] for $\delta' < \delta$. We choose $h(x) = f_{\delta'}(x) - 1$. Kapovitch showed that the inequality

$$1 \leq \frac{d(x, \hat{x})}{t} \leq \frac{1}{\cos(3\delta)}$$

holds for $x \in h^{-1}(-t)$ and $t \ll \delta' < \delta$, (see page 132 of [Kap02]). Thus, there exists a λ such that $B(\hat{x}, \frac{s}{\lambda}) \subset h^{-1}((-s, 0]) \subset B(\hat{x}, \lambda s)$ for $s \leq \frac{r}{4\lambda}$.

(2) For the convenience of readers, we add the following alternative proof of the second assertion. Let us recall that the tangent cone $(T_{\hat{x}}^-(X^n), O)$ is the Gromov-Hausdorff limit of the pointed re-scaled spaces $\{(\lambda X, \hat{x})\}_{\lambda \geq 0}$ as $\lambda \rightarrow +\infty$, i.e.

$$(\lambda X, \hat{x}) \rightarrow (T_{\hat{x}}^- X, O)$$

as $\lambda \rightarrow +\infty$, where O is the apex of the tangent cone $T_{\hat{x}}^- X$. Let $d_{O, T_{\hat{x}}^- X}(\eta) = d_{T_{\hat{x}}^- X}(O, \eta)$ and $d_{\hat{x}, \lambda X}(y) = d_{\lambda X}(y, \hat{x}) = \lambda d_X(y, \hat{x})$. We consider $f_{\lambda}(y) = \frac{1}{2}(d_{\hat{x}, \lambda X}(y, \hat{x}))^2$. By an equivalent definition of curvature ≥ -1 , $\{f_{\lambda}\}_{\lambda \geq 1}$ and f_{∞} are semi-concave functions.

Lemma 1.3.4 of [Petr07] implies that if $p_{\lambda} \rightarrow p_{\infty}$ as $\lambda \rightarrow p_{\infty}$ then $\liminf_{\lambda \rightarrow \infty} |\nabla f_{\lambda}|(p_{\lambda}) \geq |\nabla f_{\infty}|(p_{\infty})$.

Let $A_M(x, r, R) = [B_M(x, R) - B_M(x, r)]$ be an annular region. Our energy function $f_{\infty}(\eta) = \frac{1}{2}|\eta|^2$ has property $|\nabla f_{\infty}| \geq \frac{1}{8}$ on $A_{T_{\hat{x}}(X)}(0, \frac{1}{2}, 1)$. It follows from Lemma 1.3.4 of [Petr07] that, for sufficiently large

$\lambda \geq \lambda_0 > 1$, the function f_λ has no critical point on the annular region

$$A_{\lambda X}(\hat{x}, \frac{1}{2}, 1) = A_X(\hat{x}, \frac{1}{2\lambda}, \frac{1}{\lambda}).$$

Since we have

$$[B_X(\hat{x}, \frac{1}{\lambda_0}) - \{\hat{x}\}] = \cup_{\lambda \geq \lambda_0} A_X(\hat{x}, \frac{1}{2\lambda}, \frac{1}{\lambda}),$$

we conclude that the radial distance function has no critical point on the punctured ball $[B_X(\hat{x}, \varepsilon) - \{\hat{x}\}]$. $\square$

1.2. Regular values of admissible maps.

In this subsection, we recall explicit definitions of admissible mappings and their regular points introduced by Perelman.

Definition 1.8 ([Per94],[Per97] admissible maps). (1) Let X^n be a complete Alexandrov space of dimension n and $\text{curv}_{X^n} \geq c$ and $U \subset X^n$. A function $f : U \rightarrow \mathbb{R}$ is called admissible if $f(x) = \sum_{i=1}^m \phi_i(d_{A_i}(x))$ where $A_i \subset M$ is a closed subset and $\phi_i : \mathbb{R} \rightarrow \mathbb{R}$ is continuous.

(2) A map $\hat{F} : X^n \rightarrow \mathbb{R}^k$ is said to be admissible in a domain $U \subset M$ if it can be represented as $\hat{F} = G \circ F$, where $G : \mathbb{R}^k \rightarrow \mathbb{R}^k$ is bi-Lipschitz homeomorphism and each component f_i of $F = (f_1, f_2, \dots, f_k)$ is admissible.

The definition of regular points for admissible maps $\hat{F} : X^n \rightarrow \mathbb{R}^k$ on general Alexandrov spaces is rather technical. For the purpose of this paper, we only need to consider two lower dimensional cases of X^n : either X^3 is a smooth Riemannian 3-manifold or X^2 is a surface with curvature $\geq c$.

Definition 1.9 (Regular points of admissible maps for $\dim \leq 3$).

(1) Suppose that $\hat{F} : M^3 \rightarrow \mathbb{R}^2$ is an admissible map from a smooth Riemannian 3-manifold M^3 to $\mathbb{R}^2$ on a domain $U \subset M$ and $\hat{F} = G \circ F$, where $G : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is bi-Lipschitz homeomorphism and each component f_i of $F = (f_1, f_2)$ is admissible. If $\{\nabla f_1, \nabla f_2\}$ are linearly independent at $x \in U$, then x is said to be a regular point of $\hat{F}$.

(2) ([Per94] page 210). Suppose that $\hat{F} : X^2 \rightarrow \mathbb{R}^2$ is an admissible map from an Alexandrov surface X^2 of curvature $\text{curv} \geq c$ to $\mathbb{R}^2$ on a domain $U \subset X^2$ and $\hat{F} = G \circ F$, where $G : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is bi-Lipschitz homeomorphism and each component f_i of $F = (f_1, f_2)$ is admissible. Suppose that f_1 and f_2 satisfy the following conditions:

(2.a) $\langle \nabla f_1, \nabla f_2 \rangle_q \leq -\varepsilon < 0$;

(2.b) There exists $\vec{\xi} \in T_q^-(X^2)$ such that $\min\{d_q f_1(\vec{\xi}), d_q f_2(\vec{\xi})\} > \varepsilon > 0$.

Then q is called a regular point of $\hat{F}|_U$.

Remark 1.10. It is clear that Perelman's condition (2.a) implies that $\text{diam}(\Sigma_q(X^2)) > \frac{\pi}{2}$. This together with (2.b) implies that

$$\frac{\pi}{2} < \angle_q(\nabla f_1, \nabla f_2) < \pi.$$

Conversely, we would like to point out that if $\text{diam}(\Sigma_q(X^2)) > \frac{\pi}{2}$, then there exists an admissible map $F = (f_1, f_2) : U_q \rightarrow \mathbb{R}^2$ satisfying Perelman's condition (2.a) and (2.b) mentioned above, where U_q is a small neighborhood of p in X^2 . $\square$

We need to single out “bad points” (i.e., essential singularities) for which the condition

$$\text{diam}(\Sigma_q(X^2)) > \frac{\pi}{2}$$

fails. These bad points are related to the so-called extremal subsets (or essential singularities) of an Alexandrov space with curvature $\geq c$.

Definition 1.11 (Extremal points of an Alexandrov surface). *Let X^2 be an Alexandrov surface and z be an interior point of X^2 . If the diameter of space of unit tangent directions $\Sigma_z(X^2)$ has diameter less than or equal to $\frac{\pi}{2}$, i.e.*

$$\text{diam}(\Sigma_z(X^2)) \leq \frac{\pi}{2}, \quad (1.3)$$

then z is called an extremal point of the Alexandrov surface X^2 . If $\text{diam}(\Sigma_z(X^2)) > \frac{\pi}{2}$, then we say that z is a regular point of X^2 .

A direct consequence of Theorem 0.6 (i.e., Proposition 1.7) is the regularity of sufficiently small punctured disk in an Alexandrov surface.

Corollary 1.12. *Let X^2 be an Alexandrov space of curvature ≥ -1 and δ be as in Theorem 0.6. Then each point $y \in [B_{X^2}(\hat{x}, \delta) - \{\hat{x}\}]$ in punctured disk is regular.*

We recall the Perelman-Sharafutdinov gradient semi-flows for semi-concave functions.

Definition 1.13. *A curve $\phi : [a, b] \rightarrow X$ is called an f -gradient curve if for any $t \in [a, b]$*

$$\frac{d^+ \phi}{dt} = \nabla f|_{\phi(t)}. \quad (1.4)$$

It is known that if $f : X \rightarrow \mathbb{R}$ is a semi-concave function then there exists a unique f -gradient curve $\phi : [a, +\infty) \rightarrow X$ with a given initial point $\phi(0) = p$, (cf. Prop 2.3.3 of [KPT09]). We will frequently use the following result of Perelman (cf. [Per94]) and Perelman-Petrinin (cf. [PP94]).

Proposition 1.14 (Lemma 2.4.2 of [KPT09], Lemma 1.3.4 of [Petr07]). *Let $X_\alpha \rightarrow X_\infty$ be a sequence of Alexandrov space of curvature ≥ -1 which converges to an Alexandrov space X_∞ . Suppose that $f_\alpha \rightarrow f_\infty$ where $f_\alpha : X_\alpha \rightarrow \mathbb{R}$ is a sequence of λ -concave functions and $f_\infty : X_\infty \rightarrow \mathbb{R}$. Assume that $\psi_\alpha : [0, +\infty) \rightarrow X_\alpha$ is a sequence of f_α -gradient curves with $\psi_\alpha(0) = p_\alpha \rightarrow p_\infty$ and $\psi_\infty : [0, +\infty) \rightarrow X_\infty$ be the f_∞ -gradient curve with $\psi_\infty(0) = p_\infty$. Then the following is true*

(1) *for each $t \geq 0$, we have*

$$\psi_\alpha(t) \rightarrow \psi_\infty(t)$$

as $\alpha \rightarrow \infty$;

(2) $\liminf_{\alpha \rightarrow \infty} |\nabla f_\alpha|(p_\alpha) \geq |\nabla f_\infty|(p_\infty)$. *Consequently, if $\{q_\alpha\}$ is a bounded sequence of critical points of f_α , then $\{q_\alpha\}$ has a subsequence converging to a critical point q_∞ of f_∞ .*

As we pointed out earlier, the pointed spaces $\{(\frac{1}{\varepsilon}X, x)\}$ converge to the tangent cone of X at x , i.e., $(\frac{1}{\varepsilon}X, x) \rightarrow (T_x^-(X), 0)$ as $\varepsilon \rightarrow 0$, where 0 is the origin of tangent cone.

When X^2 is an Alexandrov surface of curvature ≥ -1 , it is known that X^2 is a 2-dimensional manifold. Moreover we have the following observation.

Proposition 1.15 ([Per94]). *Let X be an Alexandrov space of curvature ≥ -1 . Suppose that $\hat{x}$ is an interior point of X . Then $B_X(\hat{x}, \varepsilon)$ is homeomorphic to $B_{T_{\hat{x}}^-X}(0, \varepsilon)$, where ε is given by Proposition 1.7. Furthermore, there exists an admissible map*

$$G : T_{\hat{x}}^-(X^2) \rightarrow \mathbb{R}^2$$

such that G is bi-Lipschitz homeomorphism and G is regular at $\vec{v} \neq 0$.

Proof. This is an established result of Perelman, (cf. [Per94], [Kap07]). We provide a short proof here only for the convenience of readers. Let us first prove that $B_X(\hat{x}, \varepsilon)$ is homeomorphic to $B_{T_{\hat{x}}^-X}(0, \varepsilon)$, where ε is given by Proposition 1.7. Recall that $\{(\frac{1}{\delta}X, \hat{x})\}$ is convergent to $(T_{\hat{x}}^-X, O)$, as $\delta \rightarrow 0$. By Perelman's stability theorem (cf. Theorem 7.11 of [Kap07]), $B_{\frac{1}{\delta}X}(\hat{x}, 1)$ is homeomorphic to $B_{T_{\hat{x}}^-X}(0, 1)$ for sufficiently small δ . Thus, $B_X(\hat{x}, \delta)$ is homeomorphic to $B_{T_{\hat{x}}^-X}(0, \delta)$.

By Proposition 1.7, the function $r_{\hat{x}}(y) = d_X(y, \hat{x})$ has no critical point in punctured ball $[B_X(\hat{x}, \varepsilon) - \{\hat{x}\}]$. Thus, we can apply Perelman's fibration theorem to the following diagram:

$$\Sigma_{\hat{x}}(X) \rightarrow A_{X^2}(\hat{x}, \delta/2, \varepsilon) \xrightarrow{r_{\hat{x}}} (\delta/2, \varepsilon).$$

Consequently, we see that $A_{X^2}(\hat{x}, \frac{\delta}{2}, \varepsilon)$ is homeomorphic to a cylinder $C = (\partial B) \times (\frac{\delta}{2}, \varepsilon)$. Furthermore, the metric sphere (∂B) is homeomorphic to $\Sigma_{\hat{x}}(X)$. It follows that the metric ball $B_X(\hat{x}, \varepsilon) \sim [B_{T_{\hat{x}}^- X}(0, \delta) \cup C]$ is homeomorphic to $B_{T_{\hat{x}}^- X}(0, \varepsilon)$, where ε is given by Proposition 1.7.

It remains to construct the desired map G . If $\theta = \frac{1}{3}\text{diam}(\Sigma_{\hat{x}}(X^2))$, then $\theta \leq \pi/3$. Let us choose six vectors $\{\vec{\xi}_1, \vec{\xi}_2, \vec{\xi}_3, \vec{\xi}_4, \vec{\xi}_5, \vec{\xi}_6\} \subset \Sigma_{\hat{x}}(X^2)$ such that $\angle(\vec{\xi}_i, \vec{\xi}_j) = \theta$ for $i = j + 1$ or $\{i = 1, j = 6\}$. It is easy to construct affine map H from an Euclidean sector of angle θ to an Euclidean sector of angle $\pi/3$. In fact we can first isometrically embed an Euclidean sector as $\{(x, y) | x \geq 0, y \geq 0, 0 < \varepsilon_1 \leq \arctan(\frac{y}{x}) \leq \theta + \varepsilon_1\} \subset \mathbb{R}^2$. We could choose $h_1(x, y) = x, h_2(x, y) = \lambda y$, (see Figure 5).

Each affine map $H = (h_1, h_2)$ has height functions (distance functions from axes) up to scaling factors as its components. We can arrange the Euclidean sectors in an appropriate order so that H is admissible. For instance, we could change the role of h_1 and h_2 for adjunct sector such that, by gluing six Euclidean sectors together, we can recover $T_{\hat{x}}(X^2)$ and construct an admissible map $G : T_{\hat{x}}(X^2) \rightarrow \mathbb{R}^2$. $\square$




FIGURE 5. The construction of a regular map form $S_i \cap A_{X^2}(x_\infty, \delta, \varepsilon)$.

Recall that $\lim_{\lambda \rightarrow \infty} (\lambda X, x) = (T_x^-(X), O)$. By lifting the admissible map $G : T_x(X^2) \rightarrow \mathbb{R}^2$ to $F : \lambda X^2 \rightarrow \mathbb{R}^2$, we have the following result.

Corollary 1.16. *Let X^2 be an Alexandrov surface of curvature ≥ -1 and $\hat{x}$ be an interior point. Then there exist sufficiently small $\varepsilon > \delta > 0$ and admissible maps*

$$F_{\infty, i} : X^2 \rightarrow \mathbb{R}^2$$

such that $F_{\infty, i}$ is regular on $S_i \cap A_{X^2}(\hat{x}, \delta, \varepsilon)$ for $i = 1, 2, \dots, 6$, where $S_i \subset X^2$ is the lift of the Euclidean sector bounded by $\{\vec{\xi}_i, \vec{\xi}_{i+1}\}$ from $T_{\hat{x}}^- X^2$ to X^2 .

Proof. Let $\{h_1, h_2\}$ and $\{\vec{\xi}_1, \vec{\xi}_2, \vec{\xi}_3, \vec{\xi}_4, \vec{\xi}_5, \vec{\xi}_6\} \subset \Sigma_{\hat{x}}(X^2)$ be as in the proof of Proposition 1.15. We choose an 1-parameter family of length-minimizing geodesic segments $\hat{\gamma}_{i, \lambda} : [0, 2] \rightarrow \lambda X^2$ such that $\{\hat{\gamma}_{i, \lambda}\}$ are

converging to ξ_i in $T_{\hat{x}}(X^2)$, as $\lambda \rightarrow \infty$. We also choose another length-minimizing geodesic segment $\tilde{\gamma}_{i,\lambda} : [0, 2\delta] \rightarrow \lambda X^2$ outside the geodesic hinge $\{\hat{\gamma}_{i,\lambda}, \hat{\gamma}_{i+1,\lambda}\}$ such that $0 < \varepsilon_1/2 \leq \angle_{\hat{x}}(\tilde{\gamma}'_{i,\lambda}(0), \hat{\gamma}'_{i,\lambda}(0)) \leq \varepsilon_1$. (see Figure 5 above). We consider lifting distance functions as follows:

$$r_{\lambda X^2}(y) = d_{\lambda X^2}(\hat{x}, y)$$

and

$$f_{\lambda,i}(y) = d_{\lambda X^2}(\tilde{\gamma}_{i,\lambda}, y)$$

It follows from Proposition 1.14 that two functions $\{r_{\lambda X^2}, f_{\lambda,i}\}$ are regular on $S_i \cap A_{\lambda X^2}(\hat{x}, \frac{1}{2}, 1)$ for sufficiently large λ .

Choosing a sufficiently large λ_0 , we consider the map

$$F_{\infty,i}(\cdot) = (\lambda_0 d_{X^2}(\hat{x}, \cdot), \lambda_0 d_{X^2}(\tilde{\gamma}_i, \cdot))$$

It follows from Proposition 1.14 that $F_{\infty,i}$ is regular on the curved trapezoid-like region $S_i \cap A_{X^2}(\hat{x}, \delta, \varepsilon)$, (see Figure 5). $\square$

Recall that there is a sequence $\{M_\alpha^3\}$ convergent to X_∞^2 . We conclude Section 1 by the following circle-fibration theorem.

Theorem 1.17. *Suppose that a sequence of pointed 3-manifolds $\{(M_\alpha^3, x_\alpha)\}$ is convergent to a 2-dimensional Alexandrov surface (X_∞, x_∞) such that x_∞ is an interior point of X_∞^2 . Suppose that*

$$F_\infty : A_{X^2}(x_\infty, \delta, \varepsilon) \rightarrow A_{\mathbb{R}^2}(0, \delta, \varepsilon)$$

is a regular map as above. Then there exist maps

$$F_\alpha : A_{(M_\alpha^3, \hat{g}^\alpha)}^3(x_\alpha, \delta, \varepsilon) \rightarrow A_{\mathbb{R}^2}(0, \delta, \varepsilon)$$

such that F_α is regular for sufficiently large α . Moreover, the map F_α gives rise to a circle fibration:

$$S^1 \hookrightarrow A_{(M_\alpha^3, \hat{g}^\alpha)}^3(x_\alpha, \delta, \varepsilon) \rightarrow A_{X^2}(x_\infty, \delta, \varepsilon)$$

for sufficiently large α .

Proof. We will use the same notations as in proofs of Proposition 1.15 and Corollary 1.16. For each $F_{\infty,i}$ constructed in Corollary 1.16, we consider the map

$$F_{\infty,i} = (d_{X^2}(\hat{x}, \cdot), d_{X^2}(\tilde{\gamma}_i, \cdot))$$

up to re-scaling factors, which is regular on $S_i \cap A_{X^2}(x_\infty, \delta, \varepsilon)$. Since $B_{M_\alpha^3}(x_\alpha, r) \rightarrow B_{X^2}(x_\infty, r)$ as $\alpha \rightarrow \infty$, we can choose geodesic segment $\tilde{\gamma}_{i,\alpha}$ in M_α^3 with $\tilde{\gamma}_{i,\alpha} \rightarrow \tilde{\gamma}_i$. Then we can define a map of $F_{i,\infty}$ by

$$F_{i,\alpha} = (d_{M_\alpha^3}(x_\alpha, \cdot), d_{M_\alpha^3}(\tilde{\gamma}_{i,\alpha}, \cdot)).$$

We further choose $S_{i,\alpha} \sim F_{i,\alpha}^{-1}(S_i^*) \subset M_\alpha^3$ such that $S_{i,\alpha} \rightarrow S_i$ as $\alpha \rightarrow \infty$, where S_i^* is an Euclidean sector of $\mathbb{R}^2$ described in the

proof of Proposition 1.15 and Corollary 1.16, (see Figure 5 and Figure 6). It follows from Proposition 1.14 that $F_{i,\alpha}$ is also regular in




FIGURE 6. A regular map from $S_{i,\alpha} \cap A^3_{(M_\alpha^3, \hat{g}^\alpha)}(x_\alpha, \delta, \varepsilon)$.

$S_{i,\alpha} \cap A^3_{(M_\alpha^3, \hat{g}^\alpha)}(x_\alpha, \delta, \varepsilon)$, for sufficiently large α . Using Perelman's fibration theorem (Theorem 1.2 above), we obtain that $F_{\alpha,i}$ defines a S^1 fibration on $S_{i,\alpha} \cap A^3_{(M_\alpha^3, \hat{g}^\alpha)}(x_\alpha, \delta, \varepsilon)$. To get a global S^1 -fibration on $A^3_{(M_\alpha^3, \hat{g}^\alpha)}(x_\alpha, \delta, \varepsilon)$, we need to glue these local fibration structures together. We will discuss the detail of the gluing procedure in Section 6 below. $\square$

2. EXCEPTIONAL ORBITS, CHEEGER-GROMOLL-PERELMAN SOUL THEORY AND PERELMAN-SHARAFUTDINOV FLOWS

We begin with an example of collapsing manifolds with exceptional orbits of a circle action on a solid tori.

Example 2.0 ([CG72]). Let $\mathbb{R} \times D^2 = \{(x, y, z) | y^2 + z^2 \leq 1\}$ be an infinite long 3-dimensional cylinder. We consider an isometry $\psi_{\varepsilon, m_0} : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ given by

$$\psi_{\varepsilon, m_0} : \begin{pmatrix} x \\ y \\ z \end{pmatrix} \rightarrow \begin{pmatrix} \varepsilon \\ 0 \\ 0 \end{pmatrix} + \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos \frac{2\pi}{m_0} & -\sin \frac{2\pi}{m_0} \\ 0 & \sin \frac{2\pi}{m_0} & \cos \frac{2\pi}{m_0} \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix}$$

where m_0 is a fixed integer ≥ 2 . Let $\Gamma_{\varepsilon, m_0} = \langle \psi_{\varepsilon, m_0} \rangle$ be a sub-group generated by ψ_{ε, m_0} . It is clear that the following equation

$$\psi_{\varepsilon, m_0}^{m_0} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} x + m_0 \varepsilon \\ y \\ z \end{pmatrix}$$

holds. The quotient space $M_{\varepsilon, m_0}^3 = (\mathbb{R} \times D)/\Gamma_{\varepsilon, m_0}$ is a solid torus. Let

$$G_{\varepsilon, m_0} : \mathbb{R} \times D^2 \rightarrow M_{\varepsilon, m_0}^3$$

be the corresponding quotient map. The orbit $\mathcal{O}_{0, \varepsilon} = G_{\varepsilon, m_0}(\mathbb{R} \times \{0\})$ is an exceptional orbit in M_{ε, m_0}^3 . It is clear that such an exceptional

orbit $\mathcal{O}_{0,\varepsilon}$ has non-zero Euler number. Let $\varepsilon \rightarrow 0$, the solid tori M_{ε,m_0}^3 is convergent to $X^2 = D^2/\mathbb{Z}_{m_0}$, where $\mathbb{Z}_{m_0} = \langle \hat{\psi}_{m_0} \rangle$ is a subgroup generated by $\hat{\psi}_{m_0} : \begin{pmatrix} y \\ z \end{pmatrix} \rightarrow \begin{pmatrix} \cos \frac{2\pi}{m_0} & -\sin \frac{2\pi}{m_0} \\ \sin \frac{2\pi}{m_0} & \cos \frac{2\pi}{m_0} \end{pmatrix} \begin{pmatrix} y \\ z \end{pmatrix}$.

Let us now return to the diagram constructed in the previous section:

$$\begin{array}{ccc} A_{(M_\alpha^3, \hat{g}^\alpha)}^3(x_\alpha, \delta, \varepsilon) & \xrightarrow{F_\alpha} & \mathbb{R}^2 \\ \downarrow \text{G-H} & & \uparrow \\ A_{X^2}(x_\infty, \delta, \varepsilon) & \xrightarrow{F_\infty} & \mathbb{R}^2 \end{array}$$

Among other things, we shall derive the following theorem.

Theorem 2.1 (Solid tori around exceptional circle orbits). *Let F_α , F_∞ , $A_{(M_\alpha^3, \hat{g}^\alpha)}^3(x_\alpha, \delta, \varepsilon)$ and $A_{X^2}(x_\infty, \delta, \varepsilon)$ as in Section 1 and the diagram above. Then there is a $\delta^* > 0$ such that $B_{M_\alpha^3}(x_\alpha, \delta^*)$ is homeomorphic to a solid torus for sufficiently large α . Moreover, a finite normal cover of $B_{M_\alpha^3}(x_\alpha, \delta^*)$ admits a free circle action.*

We will establish Theorem 2.1 by using the Cheeger-Gromoll-Perelman's soul theory for *singular metrics* on open 3-dimensional manifolds with non-negative curvature, (comparing with [SY00]). It will take several steps.

2.1. A scaling argument and critical points in collapsed regions.

We start with the following observation.

Proposition 2.2 (cf. [SY00], [Yama09]). *Let $\{(B_{(M_\alpha^3, \hat{g}^\alpha)}(x_\alpha, \varepsilon), x_\alpha)\}$ be a sequence of metric balls convergent to $(B_{X^2}(x_\infty, \varepsilon), x_\infty)$ as above. Then, there is another sequence of points $\{x'_\alpha\}$ such that $x'_\alpha \rightarrow x_\infty$ as $M_\alpha^3 \rightarrow X^2$ and for sufficiently large α , the following is true:*

- (1) $\partial B_{(M_\alpha^3, \hat{g}^\alpha)}(x'_\alpha, \varepsilon)$ is homeomorphic to a quotient of torus T^2 ;
- (2) There exists $0 < \delta_\alpha < \varepsilon$ such that there is no critical point of $r_{x'_\alpha}(z) = d(z, x'_\alpha)$ for $z \in A_{(M_\alpha^3, \hat{g}^\alpha)}(x'_\alpha, \delta_\alpha, \varepsilon)$;
- (3) There is the furthest critical point z_α of the distance function $r_{x'_\alpha}$ in $B_{(M_\alpha^3, \hat{g}^\alpha)}(x'_\alpha, \delta_\alpha)$ with $\lambda_\alpha = r_{x'_\alpha}(z_\alpha) \rightarrow 0$ as $\alpha \rightarrow \infty$.

Proof. This result can be found in Shioya-Yamaguchi's paper [SY00]. For convenience of readers, we reproduce a proof inspired by Perelman and Yamaguchi (cf. [Per94], [Yama09]) with appropriate modifications.

Our choices of the desired points $\{x'_\alpha\}$ are related to certain averaging distance functions $\{h_\alpha\}$ described below. Let us first construct the limit function h_∞ of the sequence $\{h_\alpha\}$. Similar constructions related to h_∞ can be found in the work of Perelman, Grove and others (see [Per94] page 211, [PP94] page 223, [GW97] page 210, [Kap02], [Yama09]). Let $\uparrow_p^A$ denote the set of directions of geodesics from p to A in Σ_p . It is clear that if we choose $Y^2 = T_{x_\infty}(X^2)$ and if $o = o_{x_\infty}$ is the apex of $T_{x_\infty}(X^2)$, then $\uparrow_o^{\partial B_Y(o,R)} = \Sigma_o(Y^2)$ for any $R > 0$. Recall that $(\lambda X^2, x_\infty) \rightarrow (Y^2, o_{x_\infty})$ as $\lambda \rightarrow \infty$. Suppose that X^2 has curvature ≥ -1 . Applying Proposition 1.14, we see that, for any $\delta > 0$, there is a sufficiently small r such that

(2.2.4) *The minimal set $\uparrow_{x_\infty}^{\partial B_{X^2}(x_\infty, 2r)}$ is δ -dense in $\Sigma_{x_\infty}(X^2)$.*

We now choose $\delta' = \frac{\delta}{800}$ and $\delta = \frac{2\pi}{1000}$. Let $\{q_i\}_{1 \leq i \leq m}$ be a maximal (δr) -separated subset in $\partial B_{X^2}(x_\infty, r)$ and $\{q_{i,j}\}_{1 \leq j \leq N_i}$ be a $(\delta' r)$ -net in $B_{X^2}(q_i, \delta r) \cap \partial B_{X^2}(x_\infty, r)$. Yamaguchi ([Yama09]) considered

$$f_i(x) = \frac{1}{N_i} \sum_{j=1}^{N_i} d(x, q_{i,j})$$

for $i = 1, \dots, m$. Our choice of h_∞ is given by

$$h_\infty(x) = \min_{1 \leq i \leq m} \{f_i(x)\}.$$

We now verify that h_∞ has a unique maximal point x_∞ . It is sufficient to establish $h_\infty(x) \leq [r - \frac{1}{2}d(x, x_\infty)]$ for all $x \in B_{X^2}(x_\infty, \frac{r}{2})$. This can be done as follows. For each $x \in B_{X^2}(x_\infty, \frac{r}{2}) - \{x_\infty\}$, we choose i_0 such that $\angle_{x_\infty}(x, q_{i_0,j}) < 5\delta$ for $j = 1, \dots, N_{i_0}$. Suppose that $\sigma : [0, d] \rightarrow X^2$ is a length-minimizing geodesic segment of unit speed from x_∞ to x . By comparison triangle comparison theorems, one can show that if $0 < t < d(x_\infty, x)$ then $\angle_{\sigma(t)}(q_{i_0,j}, \sigma'(t)) < \frac{\pi}{4}$ for $j = 1, \dots, N_{i_0}$. It follows from the first variational formula that

$$d(\sigma(t), q_{i_0,j}) \leq r - (\cos \frac{\pi}{4})t,$$

for $j = 1, \dots, N_{i_0}$. In fact, Kapovitch [Kap02] observed that $d(\sigma(t), q_{i_0,j}) \leq [r - t \cos(3\delta)]$, (see [Kap02] page 129 or [GW97]). It follows that $h_\infty(x)_{B_{X^2}(x_\infty, \frac{r}{2})}$ has the unique maximum point x_∞ with $h_\infty(x_\infty) = r$, because $h_\infty(x) = \min_{1 \leq i \leq m} \{f_i(x)\} \leq [r - \frac{1}{2}d(x, x_\infty)]$ for $x \in B_{X^2}(x_\infty, \frac{r}{2})$.

Since $B_{M_\alpha^3}(x_\alpha, r) \rightarrow B_{X^2}(x_\infty, r)$ as $\alpha \rightarrow \infty$, we can construct a μ'_α -approximation of $\{q_{i,j,\alpha}\} \subset M_\alpha^3$ of $\{q_{i,j}\} \subset X^2$ with $\mu'_\alpha \rightarrow 0$. Let

$$f_{i,\alpha}(y) = \frac{1}{N_i} \sum_{j=1}^{N_i} d(y, q_{i,j,\alpha}) \quad \text{and} \quad h_\alpha(y) = \min_{1 \leq i \leq m} \{f_{i,\alpha}(y)\}.$$

Let A_α be a local maximum set of $h_\alpha|_{B_{M_\alpha^3}(x_\alpha, \frac{r}{4})}$ and $x'_\alpha \in A_\alpha$. Applying Proposition 1.14 to the sequence $\{h_\alpha\} \rightarrow h_\infty$, we see that $\text{diam}(A_\alpha) \rightarrow 0$ and $x'_\alpha \rightarrow x_\infty$, as $B_{M_\alpha^3}(x_\alpha, r) \rightarrow B_{X^2}(x_\infty, r)$ and $\alpha \rightarrow \infty$.

Let $r_\alpha(y) = d(y, x'_\alpha)$. Using Proposition 1.14 again, we can show that there exists a sequence $\delta_\alpha \rightarrow 0$ such that neither the function h_α nor r_α has any critical points in the annular region $A_{(M_\alpha^3, \hat{g}^\alpha)}(x'_\alpha, \delta_\alpha, \varepsilon)$, as $B_{M_\alpha^3}(x'_\alpha, r) \rightarrow B_{X^2}(x_\infty, r)$. It follows from Theorem 1.17 that the boundary $\partial B_{(M_\alpha^3, \hat{g}^\alpha)}(x'_\alpha, t)$ is homeomorphic to T^2 or Klein bottle $T^2/\mathbb{Z}^2$ for $t > \delta_\alpha$. However, $(M_\alpha^3, \hat{g}^\alpha)$ is a Riemannian 3-manifold $\partial B_{(M_\alpha^3, \hat{g}^\alpha)}(x'_\alpha, s)$ is homeomorphic to a 2-sphere for s less than the injectivity radius η_α at x'_α . Thus, we let $\mu_\alpha = \min\{\mu'_\alpha, \eta_\alpha\}$ and

$$\lambda_\alpha = \max\{d(x'_\alpha, z_\alpha) | z_\alpha \text{ is a critical point of } r_{x'_\alpha} \text{ in } B_{(M_\alpha^3, \hat{g}^\alpha)}(x'_\alpha, \delta_\alpha)\}.$$

Clearly, we have $\lambda_\alpha \in [\mu_\alpha, \delta_\alpha]$. As $\alpha \rightarrow +\infty$, we have $\lambda_\alpha \rightarrow 0$. This completes the proof of Proposition 2.2. $\square$

In what follows, we re-choose $x_\alpha = x'_\alpha$ as in the proof of Proposition 2.2 for each α . We now would like to study the sequence of re-scaled metrics

$$\tilde{g}^\alpha = \frac{1}{\lambda_\alpha^2} \hat{g}^\alpha. \quad (2.1)$$

Clearly, the curvature of $\tilde{g}^\alpha$ satisfies $\text{curv}_{\tilde{g}^\alpha} \geq -\lambda_\alpha^2 \rightarrow 0$, as $\alpha \rightarrow \infty$. By passing to a subsequence, we may assume that the pointed Riemannian 3-manifolds $\{((M_\alpha^3, \tilde{g}^\alpha), x'_\alpha)\}$ converge to a pointed Alexandrov space (Y_∞, y_∞) with non-negative curvature.

Proposition 2.3 (Lemma 3.6 of [SY00], Theorem 3.2 of [Yama09]). *Let $\{((M_\alpha^3, \tilde{g}^\alpha), x'_\alpha)\}$, X^2 , $\{\delta_\alpha\}$, $\{\lambda_\alpha\}$ and (Y_∞, y_∞) be as above. Then Y_∞ is a complete, non-compact Alexandrov space of non-negative curvature. Furthermore, we have*

- (1) $\dim Y_\infty = 3$;
- (2) Y_∞ has no boundary.

Proof. This is an established result of [SY00] and [Yama09]. We outline a proof here only for convenience of readers, using our proofs of Proposition 2.2 above and Theorem 2.4 and Proposition 2.5 below. The metric $\tilde{g}^\alpha = \frac{1}{\lambda_\alpha^2} \hat{g}^\alpha$ defined above has curvature $\geq -\lambda_\alpha^2 \rightarrow 0$, as $\alpha \rightarrow +\infty$. By our construction, the diameter of Y_∞ is infinite. Moreover, our Alexandrov space Y_∞ has no finite boundary.

It remains to show that $\dim Y_\infty = 3$. Suppose contrary, $\dim(Y_\infty^s) = 2$. Then, for each $\vec{v}^* \in \Sigma_{y_\infty}^1(Y_\infty^2)$, the subset

$$\Lambda_{\vec{v}^*}^\perp = \{\vec{w} \in \Sigma_{y_\infty}^1(Y_\infty^2) | \angle_{y_\infty}(\vec{w}, \vec{v}^*) = \frac{\pi}{2}\}$$

has at most two elements. We will find a vector $\bar{v}^*$ such that $\Lambda_{\bar{v}^*}^\perp$ has at least $N_{i_0} \geq 100$ elements for some i_0 , a contradiction to $\dim(Y_\infty) = 2$.

Our choice of $\bar{v}^*$ will be related to a tangent vector to the minimum set A of a convex function $f : Y_\infty^s \rightarrow \mathbb{R}$, which we now describe. We will retain the same notations as in the proof of Proposition 2.2 above. Let $r_{x'_\alpha}(z) = d_{M_\alpha^3}(z, x'_\alpha)$, z_α be a critical point of $r_{x'_\alpha}|_{B_{M_\alpha^3}(x'_\alpha, \frac{r}{4})}$ be as above and $\bar{z}_\alpha$ be its image in the scaled manifold $\frac{1}{\lambda_\alpha} M_\alpha^3$. Suppose that q_∞ is the limit point of a subsequence of $\{\bar{z}_\alpha\}$. It follows from Proposition 1.14 that the limiting point q_∞ must be a critical point of the distance function r_{y_∞} with $r_{y_\infty}(q_\infty) = 1$, where $r_{y_\infty}(z) = d_{Y_\infty}(z, y_\infty)$.

In addition, there exist $(N_1 + \dots + N_m)$ -many geodesic segments $\{\tilde{\gamma}_{i,j,\alpha}\}_{1 \leq j \leq N_i, 1 \leq i \leq m}$ in M_α^3 from x'_α to $q_{i,j,\alpha}$, where $\tilde{\gamma}_{i,j,\alpha} \rightarrow \tilde{\gamma}_{i,j,\infty}$ as $B_{M_\alpha^3}(x'_\alpha, r) \rightarrow B_{X^2}(x_\infty, r)$ with $\alpha \rightarrow \infty$. Let $\bar{\gamma}_{i,j,\alpha} : [0, \frac{r}{2\lambda_\alpha}] \rightarrow \frac{1}{\lambda_\alpha} M_\alpha^3$ be the re-scaled geodesic with starting point $\bar{x}'_\alpha$ in the re-scaled manifold, for $1 \leq j \leq N_i, 1 \leq i \leq m$. It can be shown that $\bar{\gamma}_{i,j,\alpha} \rightarrow \bar{\gamma}_{i,j,\infty}$ as $\frac{1}{\lambda_\alpha} M_\alpha^3 \rightarrow Y_\infty$, after passing to appropriate subsequences of $\{\alpha\}$. Therefore, we have $(N_1 + \dots + N_m)$ -many distinct geodesic rays starting from y_∞ in Y_∞ . Let us now consider limiting Busemann functions:

$$\tilde{h}_{i,j}(y) = \lim_{t \rightarrow \infty} [d(y, \bar{\gamma}_{i,j,\infty}(t)) - t] \quad \text{and} \quad \hat{h}_i(y) = \frac{1}{N_i} \sum_{j=1}^{N_i} \tilde{h}_{i,j}(y).$$

Since Y_∞ has non-negative curvature, each Busemann function $(-\tilde{h}_{i,j})$ is a convex function, (see [CG72], [Wu79] or Theorem 2.4 below). If

$$\hat{h}(y) = \min_{1 \leq i \leq m} \{\hat{h}_i\} \quad \text{and} \quad f(y) = -\hat{h}(y),$$

then f is convex. Choose $\tilde{h}_{i,j,\alpha}(x) = [\bar{d}(x, \bar{q}_{i,j,\alpha}) - \bar{d}(\bar{x}'_\alpha, \bar{q}_{i,j,\alpha})]$, $\hat{h}_{i,\alpha}(x) = \frac{1}{N_i} \sum_{j=1}^{N_i} \tilde{h}_{i,j,\alpha}(x)$ and $\hat{h}_\alpha(x) = \min_{1 \leq i \leq m} \{\hat{h}_{i,\alpha}(x)\}$ defined on $B_{\frac{1}{\lambda_\alpha} M_\alpha^3}(\bar{x}'_\alpha, \frac{r}{4\lambda_\alpha})$.

It follows that $\hat{h} = \lim_{\alpha \rightarrow \infty} \hat{h}_\alpha$. Because $\bar{x}'_\alpha$ is a maximum point of $\hat{h}_\alpha$ with $\hat{h}_\alpha(\bar{x}'_\alpha) = 0$ and $\bar{x}'_\alpha \rightarrow y_\infty$ as $\alpha \rightarrow \infty$, the point y_∞ is a critical point of the limiting function $\hat{h} = \lim_{\alpha \rightarrow \infty} \hat{h}_\alpha$ with $\hat{h}(y_\infty) = 0$.

Thus, $0 = \hat{h}(y_\infty)$ is a critical value of the *convex* function $f = (-\hat{h})$ with $\inf_{y \in Y_\infty^s} \{f(y)\} = 0$. There are two cases for $A = f^{-1}(0)$.

Case 1. If $A = \{y_\infty\}$, then it is known (cf. Proposition 2.5 below) that the distance function r_{y_∞} does not have any critical point in $[Y_\infty - \{y_\infty\}]$, which contracts to the existence of critical points q_∞ of r_{y_∞} mentioned above. Thus, this case can not happen.

Case 2. $\dim(A) \geq 1$. In this case, our proof becomes more involved. If $q_\infty \notin A = f^{-1}(0)$, then $f(q_\infty) = a > a_0 = 0$. Using the proof of

Proposition 2.5 below, we see that q_∞ can *not* be a critical point of $r_{y_\infty}(z) = d(z, y_\infty)$ either, a contradiction. Thus, $q_\infty \in A$ holds.

If, for any quasi-geodesic segment $\sigma : [a, b] \rightarrow Y_\infty$ with ending points $\{\sigma(a), \sigma(b)\} \subset \Omega$, the inclusion relation $\sigma([a, b]) \subset \Omega$ holds, then Ω is called a totally convex subset of Y_∞ . It follows from the proof of Proposition 2.5 below that the sub-level set $f^{-1}((-\infty, a])$ is totally convex. Let us choose $\vec{v}^* \in \uparrow_{y_\infty}^{q_\infty}$, where $\uparrow_p^x$ denotes the set of directions of geodesics from p to x in Σ_p . Because $\{y_\infty, q_\infty\}$ are contained in the totally convex minimal set $A = f^{-1}(0)$ of a convex function f , one has

$$\angle_{y_\infty}(\vec{v}^*, \bar{\gamma}'_{i,j,\infty}(0)) \geq \frac{\pi}{2}$$

holds for all i, j by our construction of $f = -\hat{h}$, because $\bar{\gamma}'_{i,j,\infty}(0) = \vec{s}_{i,j}$ is a support vector of $d_{y_\infty}(-f)$, (see the proof of Proposition 2.5 below). Let $\sigma : [0, \ell] \rightarrow Y$ be a geodesic segment from y_∞ to q_∞ . Since A is totally convex and f is convex, we have $\sigma([0, \ell]) \subset A$ and $f(\sigma(t)) = 0$ for all $t \in [0, \ell]$. Hence, $\hat{h}(\sigma(t)) = 0$ for all $t \in [0, \ell]$.

Recall that $\hat{h}(z) = \min_{1 \leq i \leq m} \{\hat{h}_i(z)\}$. We choose i_0 such that $\hat{h}_{i_0}(\sigma(\frac{\ell}{2})) = \min_{1 \leq i \leq m} \{\hat{h}_i(\sigma(\frac{\ell}{2}))\} = \hat{h}(\sigma(\frac{\ell}{2})) = 0$. Because $\hat{h}_{i_0}(\sigma(t))$ is a concave function of t with $0 = \hat{h}(\sigma(0)) = \hat{h}(\sigma(\ell)) \leq \min\{\hat{h}_{i_0}(\sigma(0)), \hat{h}_{i_0}(\sigma(\ell))\}$ and $\hat{h}_{i_0}(\sigma(\frac{\ell}{2})) = 0$, one concludes that $\hat{h}_{i_0}(\sigma(t)) = 0$ for all $t \in [0, \ell]$. Since $\hat{h}_{i_0}(\sigma(t)) = 0$ for all $t \in [0, \ell]$, choosing $\vec{v}^* = \sigma'(0)$ one has

$$0 = (d_{y_\infty} \hat{h}_{i_0})(\vec{v}^*) = \frac{1}{N_{i_0}} \sum_{j=1}^{N_{i_0}} [-\cos(\angle_{y_\infty}(\vec{v}^*, \bar{\gamma}'_{i_0,j,\infty}(0)))],$$

This together with inequalities $\angle_{y_\infty}(\vec{v}^*, \bar{\gamma}'_{i_0,j,\infty}(0)) \geq \frac{\pi}{2}$ implies that

$$\angle_{y_\infty}(\vec{v}^*, \bar{\gamma}'_{i_0,j,\infty}(0)) = \frac{\pi}{2}$$

holds for $j = 1, 2, \dots, N_{i_0}$. Hence, we conclude that $\bar{\gamma}'_{i_0,j,\infty}(0) \in \Lambda_{\vec{v}^*}^\perp$, for $j = 1, 2, \dots, N_{i_0}$, where $N_{i_0} \geq 100$. Therefore, we demonstrated that $\#|\Lambda_{\vec{v}^*}^\perp| \geq N_{i_0} \geq 100$. This contradicts to $\#|\Lambda_{\vec{v}^*}^\perp| \leq 2$ when $\dim(Y_\infty) = 2$. This completes the proof of the assertion $\dim(Y_\infty) = 3$. $\square$

2.2. The classification of non-negatively curved surfaces and 3-dimensional soul theory.

In what follows, if X is an open Alexandrov space of non-negative curvature, then we let $X(\infty)$ be the boundary (or called the ideal boundary) of X at infinity. For more information about the ideal boundary $X(\infty)$, one can consult with work of Shioya, (cf. [Shio94]).

In this sub-section, we briefly review the soul theory for non-negatively curved space Y_∞^k of dimension ≤ 3 . The soul theory and the splitting theorem are two important tools in the study of low dimensional collapsing manifolds.

Let X be an n -dimensional non-negatively curved Alexandrov space. Suppose that X is a non-compact complete space and that X has no boundary. Fix a point $x_0 \in X$, we consider the Cheeger-Gromoll type function

$$f(x) = \lim_{t \rightarrow +\infty} [t - d(x, \partial B(x_0, t))].$$

Let us consider the sub-level sets $\Omega_c = f^{-1}((-\infty, c])$. We will show that Ω_c is a totally convex subset for any c in Theorem 2.4 below.

H.Wu [Wu79] and Z. Shen [Shen96] further observed that

$$f(x) = \sup_{\sigma \in \Lambda} \{h_\sigma(x)\} \quad (2.2)$$

where $\Lambda = \{\sigma : [0, +\infty) \rightarrow X | \sigma(0) = x_0, d(\sigma(s), \sigma(t)) = |s - t|\}$ and h_σ is a Busemann function associated with a ray σ by

$$h_\sigma(x) = \lim_{t \rightarrow \infty} [t - d(x, \sigma(t))]. \quad (2.3)$$

Since $\Omega_c = f^{-1}((-\infty, c])$ is convex, by (2.2) we see that Ω_c contains no geodesic ray starting from x_0 . Choose $\hat{c} = \max\{c, f(x_0)\}$. Since $\Omega_{\hat{c}}$ is totally convex and contains no geodesic rays, $\Omega_{\hat{c}}$ must be compact. It follows that $\Omega_c \subset \Omega_{\hat{c}}$ is compact as well. Thus the Cheeger-Gromoll function $f(x)$ has a lower bounded

$$a_0 = \inf_{x \in M^n} \{f(x)\} = \inf_{x \in \Omega_0} \{f(x)\} > -\infty.$$

If $\Omega_{a_0} = f^{-1}(a_0)$ is a space without boundary, Ω_{a_0} is called a soul of X . Otherwise, $\partial\Omega_{a_0} \neq \emptyset$ we further consider

$$\Omega_{a_0-\varepsilon} = \{x \in \Omega_{a_0} | d(x, \partial\Omega_{a_0}) \geq \varepsilon\}$$

When X is a smooth Riemannian manifold of non-negative curvature, Cheeger-Gromoll [CG72] showed that $\Omega_{a_0-\varepsilon}$ remains to be convex. For more general case when X is an Alexandrov space of non-negative curvature, Perelman [Per91] also showed that $\Omega_{a_0-\varepsilon}$ remains to be convex, (see [Petr07] and [CDM09] as well).

Let $l_1 = \max_{x \in \Omega_{a_0}} \{d(x, \partial\Omega_{a_0})\}$ and $a_1 = a_0 - l_1$. If $\Omega_{a_1} = \Omega_{a_0-l_1}$ has no boundary, then we call Ω_{a_1} a soul of X . Otherwise, we repeat above procedure by setting

$$\Omega_{a_1-\varepsilon} = \{x \in \Omega_{a_1} | d(x, \partial\Omega_{a_1}) \geq \varepsilon\}$$

for $0 \leq \varepsilon \leq l_1$. Observe that

$$n = \dim(X) > \dim(\Omega_{a_0}) > \dim(\Omega_{a_1}) > \dots$$

Because X has finite dimension, after finitely many steps we will eventually get a sequence

$$a_0 > a_1 > a_2 > \cdots > a_m$$

such that $\Omega_{a_i-s} = \{x \in \Omega_{a_i} \mid d(x, \partial\Omega_{a_i}) \geq s\}$ for $0 \leq s \leq a_i - a_{i+1}$ and $i = 0, 1, 2, \dots, m-1$. Moreover, Ω_{a_m} is a convex subset without boundary, which is called a soul of X .

A subset Ω is said to be *totally convex* in X if for any quasi-geodesic segment $\sigma : [a, b] \rightarrow X$ with endpoints $\{\sigma(a), \sigma(b)\} \subset \Omega$, we must have $\sigma([a, b]) \subset \Omega$. The definition of quasi-geodesic can be found in [Petr07].

Theorem 2.4 (Cheeger-Gromoll [CG72], Perelman [Per91], [SY00]). *Let X be an n -dimensional open complete Alexandrov space of curvature ≥ 0 , $f(x) = \lim_{t \rightarrow +\infty} [t - d(x, \partial B(\hat{x}, t))]$, $\{\Omega_s\}_{s \geq a_m}$ and $a_0 > a_1 > \cdots > a_m$ be as above. Then the following is true.*

(1) *For each $s \geq a_0$, $\Omega_s = f^{-1}((-\infty, s])$ is a totally convex and compact subset of X ;*

(2) *If $a_i \leq s < t \leq a_{i-1}$, then*

$$\Omega_s = \{x \in \Omega_t \mid d(x, \partial\Omega_t) \geq t - s\}$$

remains to be totally convex;

(3) *The soul $N^k = \Omega_{a_m}$ is a deformation retract of X via multiple-step Perelman-Sharafutdinov semi-flows, which are distance non-increasing.*


Proof. (1) For $s \geq a_0$, we would like to show that $\Omega_s = f^{-1}((-\infty, s])$ is totally convex. Suppose contrary, there were a quasi-geodesic $\sigma : [a, b] \rightarrow X$ with

$$\max\{f(\sigma(a)), f(\sigma(b))\} \leq s$$

and c with $a < c < b$ and

$$f(\sigma(c)) > s \geq \max\{f(\sigma(a)), f(\sigma(b))\}.$$

For each integer $i \gg 1$, we choose $y_i \in \partial B(x_i, i)$ such that



$$d(y_i, \sigma(c)) = d(\partial B(x_0, i), \sigma(c)).$$

Let $\alpha_i = \angle_{\sigma(c)}(y_i, \sigma(a))$ and $\beta_i = \angle_{\sigma(c)}(y_i, \sigma(b))$. Since X has non-negative curvature and $\sigma : [a, b] \rightarrow X$ is a quasi-geodesic, it is well-known ([Petr07]) that

$$\cos \alpha_i + \cos \beta_i \geq 0.$$

It follows that

$$\min\{\alpha_i, \beta_i\} \leq \frac{\pi}{2}.$$

After passing a sub-sequence and re-indexing, we may assume that

$$\beta_{i_j} \leq \frac{\pi}{2}$$

for all $j \geq 1$. By law of cosine, we have

$$[d(y_{i_j}, \sigma(b))]^2 \leq [d(\sigma(c), y_{i_j})]^2 + |b - c|^2.$$

Therefore, we have

$$\begin{aligned} f(\sigma(b)) &\geq \lim_{j \rightarrow +\infty} [i_j - d(\sigma(b), y_{i_j})] \\ &\geq \lim_{j \rightarrow +\infty} [i_j - \sqrt{[d(\sigma(c), y_{i_j})]^2 + |b - c|^2}] \\ &= \lim_{j \rightarrow +\infty} \frac{i_j^2 - [d(\sigma(c), y_{i_j})]^2 - |b - c|^2}{i_j + \sqrt{[d(\sigma(c), y_{i_j})]^2 + |b - c|^2}} \\ &= \lim_{j \rightarrow +\infty} [i_j - d(\sigma(c), y_{i_j})] + 0 \\ &= \lim_{j \rightarrow +\infty} [i_j - d(\sigma(c), \partial B(\hat{x}, i_j))] \\ &= f(\sigma(c)) \end{aligned}$$

which is contracting to

$$f(\sigma(c)) > f(\sigma(b)).$$

Hence, Ω_c is a totally convex subset of X .

(2) Perelman [Per91] showed that if Ω_c is a convex subset of X with non-empty boundary, then the distance function

$$r_{\partial\Omega_c}(x) = d(x, \partial\Omega_c)$$

is concave for $x \in \Omega_c$, (see [Petr07] and [CDM09] as well).

(3) Because our function $(-f(x))$ and $r_{\partial\Omega_{a_i}}$ are concave in Ω_{a_i} , the corresponding semi-flows are distance non-increasing, (see Chapter 6 of [Per91], section 2 of [KPT09] or [Petr07]). Using the Perelman-Sharafutdinov flow $\frac{d^+\psi}{dt} = \frac{\nabla(-f)}{|\nabla(-f)|^2} \cdot \psi(t)$, Perelman (cf. [Per91]) showed that X is contractible to Ω_{a_0} . Let $r_i : \Omega_i \rightarrow \mathbb{R}$ be the distance function $r_i(z) = d(z, \partial\Omega_{a_i})$ if $\partial\Omega_{a_i} \neq \emptyset$ for $i = 0, 1, \dots, m-1$. For the same reasons, Ω_{a_0} is contractible to Ω_{a_1} via the Perelman-Sharafutdinov flow

$\frac{d^+\psi}{dt} = \frac{\nabla r_0}{|\nabla r_0|^2} |_{\psi(t)}$. Using the m -step Perelman-Sharafutdinov flows, we can see that the soul $N^k = \Omega_{a_m}$ is a deformation retract of X . $\square$

Proposition 2.5. *Let $f(y)$ be a convex function on Y with $a_0 = \inf_{w \in Y} \{f(w)\} > -\infty$ and $A = f^{-1}(a_0)$ as in the proofs of Proposition 2.3 and Theorem 2.4 above. Suppose that Y is an open and complete Alexandrov space with non-negative curvature and $A' \subset A$ is a closed subset of A . Then the distance function $r_{A'}(y) = d(y, A')$ from A' has no critical points in the complement $[Y - A]$ of A .*

Proof. For each $y \notin A$ and $a = f(y) > a_0$, we observe that $A \subset \text{int}(\Omega_a) = f^{-1}((-\infty, a))$. Let $\sigma : [0, \ell] \rightarrow Y$ be a length-minimizing geodesic segment of unit speed from A' to y with $\sigma(0) \in A'$ and $\sigma(\ell) = y$. Since Y has no boundary, any geodesic σ can be extended to a longer quasi-geodesic of unit speed $\tilde{\sigma} : [0, \ell + \varepsilon] \rightarrow Y$, (see [Petr07]). Since f is convex, the composition of function $t \rightarrow f(\tilde{\sigma}(t))$ remains convex for any quasi-geodesics $\tilde{\sigma}(t)$ (see [Petr07]). It follows that

$$\frac{d^+(f \circ \tilde{\sigma})}{dt}(\ell) \geq \frac{a - a_0}{\ell} > 0.$$

Let us consider a minimum direction $\vec{\xi}_{\min} \in \Sigma_y(Y)$ of $d_y(-f)$ and $\vec{s} = [-d_y(-f)(\vec{\xi}_{\min})]\vec{\xi}_{\min}$, where we used the fact that

$$[-d_y(-f)(\vec{\xi}_{\min})] \geq \frac{d^+(f \circ \tilde{\sigma})}{dt}(\ell) > 0.$$

Hence we have $\vec{s} = [-d_y(-f)(\vec{\xi}_{\min})]\vec{\xi}_{\min} \neq 0$. The vector $\vec{s}$ is called a support vector of $d_y(-f)$. For any support vector $\vec{s}$, one has (cf. [Petr07] page 143) that inequality

$$d_y(-f)(\vec{u}) \leq -\langle \vec{s}, \vec{u} \rangle$$

holds for all $\vec{u} \in \Sigma_y(Y)$, where $(-f)$ is a semi-concave function. Let $\Omega_c = f^{-1}((-\infty, c])$. When f is convex, one has

$$f(\tilde{\sigma}(t)) \leq \max\{f(\tilde{\sigma}(a)), f(\tilde{\sigma}(b))\}$$

for any quasi-geodesic $\tilde{\sigma} : [a, b] \rightarrow Y$ and $t \in [a, b]$. Thus, Ω_c is totally convex. It follows that, for any direction $\vec{w} \in \uparrow_y^{A'}$, we have $\vec{w} \in T_y(\Omega_c)$. Moreover, we have

$$d_y(-f)(\vec{w}) \geq \frac{-a_0 - (-a)}{\ell} > 0$$

for all $\vec{w} \in \uparrow_y^{A'}$, since $(-f)$ is a concave function. Because $\vec{s}$ is a support vector of $d_y(-f)$, one also has

$$0 < d_y(-f)(\vec{w}) \leq -\langle \vec{s}, \vec{w} \rangle$$

holds for $\vec{w} \in \uparrow_y^{A'}$, (cf. [Petr07] page 143). Thus, we have

$$\angle_y(\vec{w}, \vec{s}) > \frac{\pi}{2}$$

for all $\vec{w} \in \uparrow_y^{A'}$. It follows that $d_y(r_{A'})(\vec{s}) > 0$ and y is not a critical point of the distance function $r_{A'}(\cdot)$, where $y \notin A = f^{-1}(a_0)$, (compare with [Grv93]). This completes the proof of Proposition 2.5. $\square$




FIGURE 7. The minimum set A of a convex function.

Using soul theory and splitting theorem, we can classify non-negatively curved surfaces with possibly singular metrics.

Theorem 2.6. *Let X^2 be an oriented, complete and open surface of non-negative curvature. Then X^2 is either homeomorphic to $\mathbb{R}^2$ or isometric to a flat cylinder.*

Proof. It is known that X^2 is a manifold. Let $N^s = \Omega_{a_m}$ be a soul of X^2 . If the soul $N^s = \Omega_{a_m}$ is a single point, then X^2 is homeomorphic to $\mathbb{R}^2$. When $N^s = \Omega_{a_m}$ has dimension 1, then $N^1 = \Omega_{a_m}$ is isometric to embedded closed geodesic $\sigma : S^1 \rightarrow X^2$, (i.e., $N^1 = \sigma(S^1)$).

Let $\tilde{X}^2$ be the universal cover of X^2 with lifted metric and $\tilde{\sigma} : \mathbb{R} \rightarrow \tilde{X}^2$ be a lift of $N^1 = \Omega_{a_m}$ in X^2 . We observe that

$$\begin{array}{ccc} \tilde{X}^2 & \xrightarrow{\tilde{P}} & \tilde{\sigma} \\ \downarrow & & \downarrow \\ X^2 & \xrightarrow{P} & N^1 \end{array}$$

Suppose that $P : X^2 \rightarrow N^1$ is the Perelman-Sharafutdinov distance non-increasing projection from open space X^2 to its soul N^1 . Such a distance non-increasing map $P : X^2 \rightarrow N^1$ can be lifted to a distance non-increasing map $\tilde{P} : \tilde{X}^2 \rightarrow \tilde{\sigma}$. Thus $\tilde{\sigma} : \mathbb{R} \rightarrow \tilde{X}^2$ is a line in an open surface $\tilde{X}^2$ of non-negative curvature. Applying the splitting

theorem, we see that $\tilde{X}^2$ is isometric to $\mathbb{R}^2$. It follows that X^2 is a flat cylinder. $\square$

Let us now turn our attention to closed surfaces of curvature.

Corollary 2.7. *Let X^2 be a closed 2-dimensional Alexandrov space of non-negative curvature. Then the following holds:*

- (1) *If the fundamental group $\pi_1(X^2)$ is finite, then X^2 is homeomorphic to S^2 or $\mathbb{RP}^2$.*
- (2) *If the fundamental group $\pi_1(X^2)$ is an infinite group, then X^2 is isometric to a flat torus or flat Klein bottle.*

Proof. After passing through to its double when needed, we may assume that X^2 is oriented.

When $|\pi_1(X^2)| < \infty$, X^2 is covered by S^2 .

When $|\pi_1(X^2)| = +\infty$ and X^2 is oriented, for a non-trivial free homotopy class of a closed curve $[\hat{\sigma}] \neq 0$ in $\pi_1(X^2)$ with $[\hat{\sigma}^n] \neq 0$ for all $n \neq 0$, we choose a length minimizing closed geodesic $\sigma : S^1 \rightarrow X^2$. Suppose that $\tilde{X}^2$ is a universal cover of X^2 and $\tilde{\sigma} : \mathbb{R} \rightarrow \tilde{X}^2$ is a lift of σ in X^2 . Then we can check that $\tilde{\sigma}$ is a geodesic line of $\tilde{X}^2$. Thus, $\tilde{X}^2$ is isometric to $\mathbb{R}^2$. It follows that X^2 is isometric to a flat torus, whenever X^2 is oriented with $|\pi_1(X^2)| = +\infty$. $\square$

Example 2.8. When X^2 is an open surface of non-negative curvature, it might happen that Ω_{a_0} is an interval. For instance, let $\hat{Y}^2 = [0, 1] \times [0, +\infty)$ be a flat half-strip in $\mathbb{R}^2$. If we take two copies of $\hat{Y}^2$ and glue them along the boundary, the resulting surface $X^2 = \text{dbl}(\hat{Y}^2)$ is homeomorphic to $\mathbb{R}^2$. A result of Petrunin implies that $X^2 = \text{dbl}(\hat{Y}^2)$ still have non-negative curvature (e.g., [Petr07] or [BBI01]). In this case, we have Ω_{a_0} is an interval. Of course, the soul $N^s = \Omega_{a_1}$ of X^2 is a single point.

We now say a few words for non-negatively curved surfaces X^2 with non-empty convex boundary. By definition of surface X^2 with curvature $\geq k$, its possibly non-empty boundary ∂X^2 must be convex.

Corollary 2.9. *Let X^2 be a surface with non-negative curvature and non-empty boundary. Then*

- (1) *If X^2 is compact, then X^2 is either homeomorphic to D^2 or isometric to $S^1 \times [0, l]$ or a flat Möbius band;*
- (2) *If X^2 is non-compact and oriented, then X^2 is either homeomorphic to $[0, +\infty) \times \mathbb{R}$ or isometric to one of three types: $S^1 \times [0, +\infty)$, a half flat strip or $[0, l] \times (-\infty, +\infty)$.*

Proof. If we take two copies of X^2 and glue them together along their boundaries, the resulting surface $\text{dbl}(X^2)$ still has curvature ≥ 0 , due to a result of Petrunin [Petr07]. Clearly, $\text{dbl}(X^2)$ has no boundary.

(1) When $\text{dbl}(X^2)$ is compact and oriented, then $\text{dbl}(X^2)$ is homeomorphic to the unit 2-sphere or is isometric to a flat strip. Hence, X^2 is either homeomorphic to D^2 or isometric to $S^1 \times [0, l]$ or a flat Möbius band.

(2) When $\text{dbl}(X^2)$ is non-compact, then $\text{dbl}(X^2)$ is homeomorphic to $\mathbb{R}^2$ or isometric to $S^1 \times \mathbb{R}$ or X^2 is isometric to $[0, \ell] \times [0, \infty)$.

To verify this assertion, we consider the soul N^s of $\text{dbl}(X^2)$. If N^s is a circle, then $\text{dbl}(X^2)$ is isometric to an infinite flat cylinder: $S^1(r) \times \mathbb{R}$. If the soul N^s is a point, then $\text{dbl}(X^2)$ is homeomorphic to $\mathbb{R}^2$.

There is a special case which we need to single out: X^2 is isometric to $[0, \ell] \times [0, \infty)$. We will elaborate this special case in Section 5 below. $\square$

Remark 2.10. In Section 5 below, we will estimate the number of extremal points, i.e. essential singularities, on surfaces with non-negative curvature, using multi-step Perelman-Sharafutdinov flows associated with the Cheeger-Gromoll convex exhaustion.

Finally, we would like to classify all non-negatively curved open 3-manifolds with possibly singular metrics.

Theorem 2.11 ([SY00]). *Let Y_∞^3 be an open complete 3-manifold with a possibly singular metric of non-negative curvature. Suppose that Y_∞^3 is oriented and N^s is a soul of Y_∞^3 . Then the following is true.*

- (1) *When $\dim(N^s) = 1$, then the soul of Y_∞^3 is isometric to a circle. Moreover, its universal cover $\tilde{Y}_\infty^3$ is isometric to $\tilde{X}^2 \times \mathbb{R}$, where $\tilde{X}^2$ is homeomorphic to $\mathbb{R}^2$;*
- (2) *When $\dim(N^s) = 2$, then the soul of Y_∞^3 is homeomorphic to S^2/Γ or T^2/Γ . Furthermore, Y_∞^3 is isometric to one of four spaces: $S^2 \times \mathbb{R}$, $\mathbb{R}P^2 \ltimes \mathbb{R} = (S^2 \times \mathbb{R})/\mathbb{Z}_2$, $T^2 \times \mathbb{R}$ or $K^2 \ltimes \mathbb{R} = (T^2 \times \mathbb{R})/\mathbb{Z}_2$, where K^2 is the flat Klein bottle and $\mathbb{R}P^2 \ltimes \mathbb{R}$ is homeomorphic to $[\mathbb{R}P^3 - \bar{B}^3(x_0, \epsilon)]$;*
- (3) *When $\dim(N^s) = 0$, then the soul of Y_∞^3 is a single point and Y_∞^3 must be homeomorphic to $\mathbb{R}^3$.*

Proof. This theorem is entirely due to Shioya-Yamaguchi [SY00]. A special case of Theorem 2.11 for smooth open 3-manifold with non-negative curvature was stated as Theorem 8.1 in Cheeger-Gromoll's paper [CG72].

Shioya-Yamaguchi's proof is quiet technical, which occupied the half of their paper [SY00]. For convenience of readers, we present an alternative shorter proof of Shioya-Yamaguchi's soul theorem for 3-manifolds with possible singular metrics.

Case 1. When the soul N^1 of Y_∞^3 is a closed geodesic σ^1 , there are distance non-increasing multi-step Perelman-Sharafutdinov retractions from Y_∞^3 to σ^1 . Thus, σ^1 is length-minimizing in its free homotopy class. It follows that the lifting geodesic $\tilde{\sigma}^1$ is a geodesic line in the universal covering space $\tilde{Y}_\infty^3$ of Y_∞^3 . Using the splitting theorem (cf. [BBI01]) for non-negatively curved space $\tilde{Y}_\infty^3$, we see that $\tilde{Y}_\infty^3$ is isometric to $\tilde{X}^2 \times \mathbb{R}$, where $\tilde{X}^2$ is a contractible surface with non-negative curvature. Hence, $\tilde{X}^2$ is homeomorphic to $\mathbb{R}^2$.

Case 2. When the soul N^2 of Y_∞^3 is a surface X^2 , we observe that $X^2 = f^{-1}(a_0)$ is a convex subspace of Y_∞^3 , where $f(x) = \lim_{t \rightarrow \infty} [t - d(x, \partial B_{Y_\infty^3}(\hat{x}, t))]$ and $a_0 = \inf_{x \in Y_\infty^3} \{f(x)\}$. Since f is convex and Y_∞^3 has non-negative curvature, X^2 has non-negative curvature as well. By Corollary 2.9, we see that X^2 is either homeomorphic to a quotient of S^2 or isometric to a quotient of a flat torus.

For this case, our strategy goes as follows. We will show that there is a “*normal line bundle*” over the soul X^2 . After passing its double cover if needed, we may assume that such a “*normal line bundle*” is topologically trivial in Y_∞^3 . In this case, with some extra efforts, one can show that there is a geodesic line $\hat{\sigma}^1$ orthogonal to X^2 in Y_∞^3 . Thus, the space Y_∞^3 (or its double cover) splits isometrically to $X^2 \times \mathbb{R}$.

Here is the detail of our “*normal line bundle*” argument. For each point x in the soul X^2 , its unit tangent space $\Sigma_x^1(X^2)$ is homeomorphic to S^1 . Recall that the space of unit tangent directions $\Sigma_{y_\infty}^2(Y_\infty^3)$ of Y_∞^3 at y is homeomorphic to the sphere S^2 , because Y_∞^3 is a 3-manifold. Observe that $\Sigma_x^1(X^2)$ is a convex subset of $\Sigma_{y_\infty}^2(Y_\infty^3)$. Moreover we see that $\Sigma_x^1(X^2)$ divides $\Sigma_{y_\infty}^2(Y_\infty^3)$ into exactly two parts:

$$[\Sigma_{y_\infty}^2(Y_\infty^3) - \Sigma_x^1(X^2)] = \Omega_{x,+}^2 \cup \Omega_{x,-}^2$$

Since the curvature of $\Sigma_{y_\infty}^2(Y_\infty^3)$ is greater than or equal to 1, using Theorem 6.1 of [Per91] (cf. [Petr07]), we obtain that there is a unique unit vector $\xi_\pm \in \Omega_{x,\pm}^2$ such that

$$\ell_\pm = \angle_x(\xi_\pm, \Sigma_x^1(X^2)) = \max\{\angle_x(w_\pm, \Sigma_x^1(X^2)) \mid w_\pm \in \Omega_{x,\pm}^2\}.$$

We claim that $\ell_\pm \leq \frac{\pi}{2}$. Suppose contrary, $\ell_\pm > \frac{\pi}{2}$ were true. We derive a contradiction as follows. Let $\psi : [0, \ell] \rightarrow \Sigma_{y_\infty}^2(Y_\infty^3)$ be a length-minimizing geodesic segment of unit speed from $\Sigma_x^1(X^2)$ of length $\ell \geq \frac{\pi}{2}$ with $\psi(0) = u \in \Sigma_x^1(X^2)$ and $d(\psi(\ell), \Sigma_x^1(X^2)) = \ell$. We now choose

another geodesic segment $\eta : [0, \delta] \rightarrow \Sigma_x^1(X^2)$ be a geodesic segment of unit speed with $\eta(0) = u$. Since $\text{curv}_{\Sigma^2} \geq 1$, applying the triangle comparison theorem (cf. [BBI01]) to the our geodesic hinge at u in $\Sigma_{y_\infty}^2(Y_\infty^3)$, we see that $d_{\Sigma^2}(\psi(\frac{\pi}{2}), \eta(\delta)) \leq \frac{\pi}{2}$. Thus, for the point $w = \psi(\frac{\pi}{2})$, there are at least two points $\{u, \eta(\delta)\} \subset \Sigma_x^1(X^2)$ with angular distance $d_{\Sigma^2}(w, u) = d_{\Sigma^2}(w, \eta(\delta)) = \frac{\pi}{2}$. In another words, there were at least two distinct length-minimizing geodesic segments from $w = \psi(\frac{\pi}{2})$ to $\Sigma_x^1(X^2)$. Hence, $\psi|_{[0, \frac{\pi}{2} + \varepsilon]}$ is no longer length-minimizing for any $\varepsilon > 0$, a contradiction. It follows that $\ell_\pm \leq \frac{\pi}{2}$. Moreover, the equality $\ell_\pm = \frac{\pi}{2}$ holds if and only if $\Sigma_{y_\infty}^2(Y_\infty^3)$ is isometric to the two point spherical suspension of $\Sigma_x^1(X^2)$. In this case, $T_x(Y_\infty^3)$ is isometric to $T_x(X^2) \times \mathbb{R}$.

Recall that $X^2 = f^{-1}(a_0)$ is a level set of the Busemann function. We can write $f(x) = [c - d(x, f^{-1}(c))]$ for $x \in f^{-1}(-\infty, c]$. By the first variational formula (cf. [BBI01] page 125), we see that

$$\ell_\pm \geq \frac{\pi}{2}.$$

Combining our earlier inequality $\ell_\pm \leq \frac{\pi}{2}$, we see that $\ell_\pm = \frac{\pi}{2}$. Therefore, we conclude that $T_x(Y_\infty^3)$ is isometric to $T_x(X^2) \times \mathbb{R}$. Hence, there is a “normal line bundle” over the soul X^2 . After passing its double cover if necessary, such a “normal line bundle” of X^2 in Y_∞^3 is topologically trivial. Thus, we assume that $[Y_\infty^3 - X^2] = \Omega_+^3 \cup \Omega_-^3$ has exactly two ends, where we replace Y_∞^3 by its double cover $\hat{Y}^3$ if needed. For each end and each $x \in X^2$, there exists a ray $\sigma_{x, \pm} : (0, \infty) \rightarrow \Omega_\pm^3$ with starting point x . One can verify that $\sigma_{x, -} \cup \sigma_{x, +}$ is a geodesic line in Y_∞^3 (or in its double cover). By the splitting theorem, we conclude that Y_∞^3 (or its double cover) is isometric to $X^2 \times \mathbb{R}$.

(3) When the soul N^k of Y_∞^3 is a single point $\{y_\infty\}$, our proof becomes more involved. Let $f(x) = \lim_{t \rightarrow \infty} [t - d(x, \partial B_{Y_\infty^3}(\hat{x}, t))]$ and $a_0 = \inf_{x \in Y_\infty^3} \{f(x)\}$ be as above. There are three possibilities for $\dim[f^{-1}(a_0)] = 0, 1, 2$.

Subcase 3.0. $\dim[f^{-1}(a_0)] = 0$ and $A = f^{-1}(a_0) = \{y_\infty\}$.

In this subcase, the space of unit tangent directions $\Sigma_{y_\infty}^2(Y_\infty^3)$ at y is homeomorphic to the sphere S^2 and its tangent cone $T_{y_\infty}(Y_\infty^3)$ is homeomorphic to $\mathbb{R}^3$. Recall that the pointed spaces $(\lambda Y_\infty^3, y_\infty)$ is convergent to the tangent cone $(T_{y_\infty}(Y_\infty^3), O)$ as $\lambda \rightarrow \infty$, where O is the origin of $T_{y_\infty}(Y_\infty^3)$.

By the pointed version of Perelman’s stability theorem (cf. Theorem 7.11 of [Kap07]), we see that for sufficiently small ε , $(B_{\frac{1}{\varepsilon} Y_\infty^3}(y_\infty, 1), y_\infty)$ is homeomorphic to $(B_{T_{y_\infty}(Y_\infty^3)}(O, 1), O)$. It follows that $B_{Y_\infty^3}(y_\infty, \varepsilon)$ is

homeomorphic to the unit ball D^3 for sufficiently small $\varepsilon > 0$, because Y_∞^3 is a 3-manifold.

We now use Perelman's fibration theorem to complete our proof for this subcase. It follows from Proposition 2.5 that r_A has no critical value in $(0, \infty)$. Perelman's fibration theorem (our Theorem 1.2 above) implies that there is a fibration structure

$$(\partial D^3) \rightarrow [Y_\infty^3 - U_\varepsilon(A)] \xrightarrow{r_A} (\frac{\varepsilon}{2}, \infty)$$

It follows that $[Y_\infty^3 - U_\varepsilon(A)]$ is homeomorphic to $S^2 \times (\frac{\varepsilon}{2}, \infty)$ and that Y_∞^3 is homeomorphic to $D^3 \cup [S^2 \times \mathbb{R}]$. Thus, Y_∞^3 is homeomorphic to $\mathbb{R}^3$, for this subcase.

Subcase 3.1. $\dim[f^{-1}(a_0)] = 1$ and $A = f^{-1}(a_0) = \sigma$ is a geodesic segment.

It follows from Proposition 2.5 that r_A has no critical value in $(0, \infty)$. For the same reasons as above, it remains important to verify that $U_\varepsilon(A)$ is homeomorphic to D^3 .

Let $\sigma : [0, \ell] \rightarrow Y^3$ be as above and $\sigma([0, \ell])$ be the minimal set of f . We denote a ε -neighborhood of A by $U_\varepsilon(A)$. Let $A_s = \sigma([s, \ell - s])$ for some $s > 0$. We observe that

$$U_\varepsilon(A_0) = B_\varepsilon(\sigma(0)) \cup U_\varepsilon(A_s) \cup B_\varepsilon(\sigma(\ell))$$

for $s > 0$. For the same reason as in Subcase 3.0 above, both $B_\varepsilon(\sigma(0))$ and $B_\varepsilon(\sigma(\ell))$ are homeomorphic to D^3 , because Y_∞^3 is a 3-manifold. It is sufficient to show that $U_\varepsilon(A_s)$ is homeomorphic to a finite cylinder $C = [s, \ell - s] \times D^2$ for sufficient small s .

Let $p = \sigma(0)$. We consider the distance function $r_p(y) = \text{d}_{Y_\infty^3}(y, p)$. We observe that the distance function has no critical point on geodesic sub-segment $\sigma([s, \ell - s])$. A result of Petrunin (cf. [Petr07] page 142) asserts that if $x_n \rightarrow x$ as $n \rightarrow \infty$, then $\liminf_{n \rightarrow \infty} |\nabla_{x_n} r_p| \geq |\nabla_x r_p|$. Hence, there exists a sufficiently small $\varepsilon > 0$ such that r_p has no critical point in $U_\varepsilon(A_s)$. For the same reason as in Subcase 3.0, we can apply Perelman's fibration theorem to our case:

$$D^2 \rightarrow U_\varepsilon(A_s) \xrightarrow{r_p} (s, \ell - s),$$

where we used the fact that $[\partial B_\varepsilon(\sigma(0))] \cap U_\varepsilon(A_{s/2})$ is homeomorphic to D^2 . It follows that $U_\varepsilon(A_s)$ is homeomorphic to a finite cylinder $C = (s, \ell - s) \times D^2$. Therefore, $U_\varepsilon(A_0)$ is homeomorphic to D^3 . It follows that $Y_\infty^3 \sim [D^3 \cup (S^2 \times \mathbb{R})]$ is homeomorphic to $\mathbb{R}^3$.

Subcase 3.2. $\dim[f^{-1}(a_0)] = 2$ and $A = f^{-1}(a_0) = \Omega_0^2 \sim D^2$ is a totally convex surface with boundary.

For the same reason as the two subcases above, it is sufficient to establish that $U_\varepsilon(A)$ is homeomorphic to the unit 3-ball D^3 :

$$U_\varepsilon(A) \sim D^3.$$

Let $A_s = \{x \in \Omega_0^2 \mid d(x, \partial\Omega_0^2) \geq s\}$. By our discussion in Case 2 above, we see that for each interior point $x \in \Omega_0^2$, there is a unique *normal line* orthogonal to Ω_0^2 at x . Thus, the interior $\text{int}(\Omega_0^2)$ has a normal line bundle $\mathbb{R}$. Because $\text{int}(\Omega_0^2)$ is contractible to a soul point y_0 , any line bundle over $\text{int}(\Omega_0^2)$ is topologically trivial.

In this subcase, our technical goals are to show the following:

(3.2a) $U_\varepsilon(A_s)$ is homeomorphic to $D^2 \times (-\varepsilon, \varepsilon)$;

(3.2b) $U_\varepsilon(\partial A_0)$ is homeomorphic to a solid tori $(\partial D^2) \times D_\varepsilon^2 = S^1 \times D_\varepsilon^2$.

To establish (3.2a), we use a theorem of Perelman (cf. Theorem 6.1 of [Per91]) to show that there is a product metric on a subset $U_\varepsilon(A_s)$ of Y_∞^3 . Inspired by Perelman, we consider the distance function $r_{A_0}(y) = d_{Y_\infty^3}(y, A_0)$. Since Y_∞^3 has non-negative curvature and $\text{int}(A_s)$ is *weakly concave* towards its complement $[U_{s/4}(A_s) - A_s]$, Perelman observed that r_{A_0} is *concave* on $[U_{s/4}(A_s) - A_s]$, (see the proof of Theorem 6.1 in [Per91], [Petr07] or [CDM09]). We already showed that for each interior point $x \in A_0$, there is a unique *normal line* orthogonal to A_0 at x . With extra efforts, we can show that, for each interior point $x \in \text{int}(A_0)$ and each unit normal direction $\xi_\pm \perp_x (\text{int} A_0)$, there is a unique ray $\sigma_{x,\pm} : [0, \infty) \rightarrow Y_\infty^3$ with $\sigma_{x,\pm}(0) = x$ and $\sigma'_{x,\pm}(0) = \xi_\pm$. Moreover, we have

$$f(\sigma_{x,\pm}(t)) = a_0 + t.$$

Therefore each $y \in [U_{s/4}(A_s) - A_s]$ with $s > 0$, we have

$$\nabla(-f)|_y = -\nabla r_{A_0}|_y.$$

Hence, our Busemann function f is both convex and concave on the subset $[U_{s/4}(A_s) - A_s]$. Thus, for any geodesic segment $\varphi : [a, b] \rightarrow [U_{s/4}(A_s) - A_s]$, the function $f(\varphi(t))$ is a linear function in t . Using the fact that $f(\varphi(t))$ is a linear function in t and the sharp version of triangle comparison theorem (cf. [BGP92]), we can show that there is a sub-domain V of Y_∞^3 such that the metric of Y_∞^3 on V splits isometrically as

$$V = \text{int}(A_0) \times \mathbb{R}.$$

Since $\text{int}(A_0)$ is homeomorphic to D^2 , we conclude that $U_\varepsilon(A_s)$ is homeomorphic to $D^2 \times (-\varepsilon, \varepsilon)$ (compare with the proof of Theorem 5.4 below). Hence, our Assertion (3.2a) holds.

It remains to verify (3.2b). We consider the doubling surface $\text{dbl}(A_0) = A_0 \cup_{\partial A_0} A_0$. It follows from a result of Petrunin that $\text{dbl}(A_0)$ has non-negative curvature. By Proposition 1.7, we see that the essential singularities (extremal points) in $\text{dbl}(A_0)$ are isolated. Thus, there are only finitely many points $\{x_1, \dots, x_k\}$ on ∂A_0 such that

$$\text{diam}[\Sigma_{x_j}(A_0)] \leq \frac{\pi}{2}$$

for $j = 1, \dots, k$. We can divide our boundary curve ∂A_0 into k -many arcs, say $[\partial A_0 - \{x_1, \dots, x_k\}] = \cup \gamma_j$. Using a similar argument as in Subcase 3.1, we can show that, for each γ_j , its ε -neighborhood $U_\varepsilon(\gamma_j)$ is homeomorphic to a finite cylinder $C_j \sim [D^2 \times (0, \ell_j)]$. Since Y_∞^3 is a 3-manifold, by the proof of Proposition 1.15, we know that $B_{Y_\infty^3}(x_j, \varepsilon)$ is homeomorphic to D^3 . Consequently, we have

$$U_\varepsilon(\partial A_0) = [\cup C_j] \bigcup [\cup B_{Y_\infty^3}(x_j, \varepsilon)],$$

which is homeomorphic to a solid tori $D^2 \times S^1$. This completes our proof of the assertion that $U_\varepsilon(A_0) \sim \{U_\varepsilon(\partial A_0) \cup [A_0 \times (-\varepsilon, \varepsilon)]\}$ is homeomorphic to D^3 . Therefore, $Y_\infty^3 \sim [D^3 \cup (S^2 \times \mathbb{R})]$ is homeomorphic to $\mathbb{R}^3$.

We now finished the proof of our soul theorem for all cases. $\square$

2.3. Applications of the soul theory to proof of Theorem 2.1.

Using Theorem 2.11, we can complete the proof of Theorem 2.1.

Proof of Theorem 2.1. Let λ_α and $\tilde{g}^\alpha$ be defined by (2.1). We may assume that $\{((M_\alpha^3, \tilde{g}^\alpha), x_\alpha)\}$ is convergent to a pointed Alexandrov space (Y_∞^3, y_∞) of non-negative curvature, by replacing x_α with x'_α in Proposition 2.2 if needed. By Proposition 2.3, we see that the limiting space Y_∞^3 is a non-compact and complete space of $\dim Y_\infty^3 = 3$. Furthermore, Y_∞^3 has no boundary. By Perelman's stability theorem (cf. [Kap07]), we see that the limit space Y_∞^3 is a topological 3-manifold.

By Theorem 1.17, we see that $\partial B_{(M_\alpha^3, \tilde{g}^\alpha)}(x_\alpha, r)$ is homeomorphic to a quotient of the 2-torus T^2 . The notion of ideal boundary $Y_\infty^3(\infty)$ of Y_∞^3 can be found in [Shio94]. In our case, the ideal boundary $Y_\infty^3(\infty)$ at infinity of Y_∞^3 is homeomorphic to a circle $\partial B_{X^2}(x_\infty, r)$. We will verify that $N^k \sim S^1$ as follows. Let $U_\varepsilon(N^k)$ be a ε -tubular neighborhood of the soul N^k in Y_∞^3 . By Perelman's stability theorem and our assumption that M_α^3 is oriented, we observe that, for sufficiently large α , the boundary $\partial U_r(N^k)$ is homoeomorphic to $\partial B_{(M_\alpha^3, \tilde{g}^\alpha)}(x_\alpha, r') \sim T^2$. Thus, the soul N^k of Y^3 must be a circle S^1 . In this case, it follows from Theorem 2.11 that the metric on Y^3 (or on its universal cover) splits. Therefore, it follows from Perelman's stability theorem (cf. [Kap07])

that $B_{(M_\alpha^3, \tilde{g}^\alpha)}(x_\alpha, r')$ is homeomorphic to a solid tori, which is foliated by orbits of a free circle action for sufficiently large α . We discuss more about gluing and perturbing our local circle actions in Section 6. $\square$

We will use Theorem 2.1-2.11 to derive more refined results for collapsing 3-manifolds with curvature ≥ -1 in upcoming sections.

3. ADMISSIBLE DECOMPOSITIONS FOR COLLAPSED 3-MANIFOLDS

Let $S^2(\varepsilon)$ be a round sphere of constant curvature $\frac{1}{\varepsilon^2}$. It is clear that $S^2(\varepsilon) \times [a, b]$ is convergent to $[a, b]$ with non-negative curvature as $\varepsilon \rightarrow 0$. The product space $W_\varepsilon = S^2(\varepsilon) \times [a, b]$ is not a graph manifold. However, if W_ε is contained in the interior of collapsed 3-manifold M_α^3 with boundary, then for topological reasons, W_α still has a chance to become a part of graph-manifold M_α^3 .

Let us now use the language of Cheeger-Gromov's F -structure theory to describe 3-dimensional graph-manifold. It is known that a 3-manifold M^3 is a graph-manifold if and only if M^3 admits an F -structure of positive rank, which we now describe.

An F -structure, $\mathcal{F}$, is a topological structure which extends the notion of torus action on a manifold, (see [CG86] and [CG90]). In fact, the more significant concept is that of atlas (of charts) for an F -structure.

An atlas for an F -structure on a manifold M^n is defined by a collection of triples $\{(U_i, V_i, T^{k_i})\}$, called charts, where $\{U_i\}$ is an open cover of M^n and the torus, T^{k_i} , acts effectively on a finite normal covering, $\pi_i : V_i \rightarrow U_i$, such that the following conditions hold:

- (3.1) There is a homomorphism, $\rho_i : \Gamma_i = \pi_1(U_i) \rightarrow \text{Aut}(T^{k_i})$, such that the action of T^{k_i} extends to an action of the semi-direct product $T^{k_i} \rtimes_{\rho_i} \Gamma_i$, where $\pi_1(U)$ is the fundamental group of U ;
- (3.2) If $U_{i_1} \cap U_{i_2} \neq \emptyset$, then $U_{i_1} \cap U_{i_2}$ is connected. If $k_{i_1} \leq k_{i_2}$, then on a suitable finite covering of $U_{i_1} \cap U_{i_2}$, their lifted tori-actions commute after appropriate re-parametrization.

The compatibility condition (3.2) on lifted actions implies that M^n decomposes as a disjoint union of orbits, $\mathcal{O}$, each of which carries a natural flat affine structure. The orbit containing $x \in M^n$ is denoted by $\mathcal{O}_x$. The dimension of an orbit of minimal dimension is called the rank of the structure.

Proposition 3.1 ([CG86], [CG90]). *A 3-dimensional manifold M^3 with possible non-empty boundary is a graph-manifold if and only if M^3 admits an F -structure of positive rank.*

For 3-dimensional manifolds, we will see that 7 out of 8 geometries admits F-structure. Therefore, there seven types of locally homogeneous spaces are graph-manifolds.

Example 3.2. Let M^3 be a closed locally homogeneous space of dimension 3, such that its universal covering spaces $\tilde{M}^3$ is isometric to seven geometries: $\mathbb{R}^3, S^3, \mathbb{H}^2 \times \mathbb{R}, S^2 \times \mathbb{R}, SL(2, \mathbb{R}), Nil$ and Sol . Then M^3 admits an F-structure and hence it is a graph-manifold.

Let us elaborate this issue in detail as follows.

- (i) If $M^3 = \mathbb{R}^3/\Gamma$ is a flat 3-manifold, then it is covered by 3-dimensional torus. Hence it is a graph-manifold.
- (ii) If $M^3 = S^3/\Gamma$ is a lens space, then its universal cover S^3 admits the classical Hopf fibration:

$$S^1 \rightarrow S^3 \rightarrow S^2.$$

It follows that M^3 is a graph-manifold.

- (iii) If $M^3 = (\mathbb{H}^2 \times \mathbb{R})/\Gamma$ is a closed 3-manifold, then a theorem of Eberlein implies that a finite normal cover $\hat{M}^3$ of M^3 is diffeomorphic to $N^2 \times S^1$, where N^2 is a closed surface of genus ≥ 1 , (see Proposition 5.11 of [CCR01], [CCR04]).
- (iv) If $M^3 = (S^2 \times \mathbb{R})/\Gamma$, then a finite cover is isometric to $S^2 \times S^1$. Clearly, M^3 is a graph-manifold. We should point out that a quotient space $(S^2 \times S^1)/\mathbb{Z}_2$ may be homeomorphic to $\mathbb{RP}^3 \# \mathbb{RP}^3$.
- (v) If $M^3 = \tilde{SL}(2, \mathbb{R})/\Gamma$, then a finite cover $\hat{M}^3$ of M^3 is diffeomorphic to the unit tangent bundle of a closed surface N_k^2 of genus $k \geq 2$. Thus, we may assume that $\hat{M}^3 = SN_k^2 = \{(x, \vec{v}) | x \in N_k^2, \vec{v} \in T_x(N_k^2), |\vec{v}| = 1\}$. Clearly, there is a circle fibration

$$S^1 \rightarrow \hat{M}^3 \rightarrow N_k^2.$$

It follows that M^3 is a graph-manifold.

- (vi) If $M^3 = Nil/\Gamma$, then the universal cover

$$\tilde{M}^3 = Nil = \left\{ \left(\begin{pmatrix} 1 & x & z \\ 0 & 1 & y \\ 0 & 0 & 1 \end{pmatrix} \right) \middle| x, y, z \in \mathbb{R} \right\}$$

is a 3-dimensional Heisenberg group. Let

$$\hat{\Gamma} = \left\{ \left(\begin{pmatrix} 1 & m & k \\ 0 & 1 & n \\ 0 & 0 & 1 \end{pmatrix} \right) \middle| k, m, n \in \mathbb{Z} \right\}$$

- be the integer lattice group of Nil . A finite cover $\hat{M}^3$ of M^3 is a circle bundle over a 2-torus. Therefore M^3 is a graph-manifold, which can be a diameter-collapsing manifold.
- (vii) If $M^3 = Sol/\Gamma$, then M^3 is foliated by tori, Möbius bands or Klein bottles, which is a graph-manifold.

Let us consider a graph-manifold which is not a compact quotient of a homogeneous space.

Example 3.3. Let N_i^2 be a surface of genus ≥ 2 and with a boundary circle, for $i = 1, 2$. Clearly, $\partial(N_i^2 \times S^1) = S^1 \times S^1$. We glue $N_1^2 \times S^1$ to $S^1 \times N_2^2$ along their boundaries with S^1 -factor switched. The resulting manifold $M^3 = (N_1^2 \times S^1) \cup (S^1 \times N_2^2)$ does not admit a global circle fibration, but M^3 is a graph-manifold.

As we pointed out above, $S^2 \times [a, b]$ can be collapsed to an interval $[a, b]$ with non-negative curvature. Suppose that W is a portion of collapsed 3-manifold M^3 such that W is diffeomorphic to $S^2 \times [a, b]$. We need to glue extra solid handles to W so that our collapsed 3-manifold under consideration becomes a graph-manifold. For this purpose, we divided $S^2 \times [a, b]$ into three parts. In fact, S^2 with two disks removed, $S^2 - (D_1^2 \sqcup D_2^2)$, is diffeomorphic to an annulus A . Thus, $S^2 \times [a, b]$ has a decomposition

$$S^2 \times [a, b] = (D_1^2 \times [a, b]) \sqcup (D_2^2 \times [a, b]) \sqcup (A \times [a, b]).$$

The product space $A \times [a, b]$ clearly admits a free circle action, and hence is a graph-manifold. For solid cylinder part $D_i^2 \times [a, b]$, if one can glue two solid cylinders together, then one might end up with a solid torus $D^2 \times S^1$ which is again a graph-manifold.

We will decompose a collapsed 3-manifold M_α^3 with curvature ≥ -1 into four major parts according to the dimension k of limiting set X^k :

$$M_\alpha^3 = V_{\alpha, X^0} \cup V_{\alpha, \text{int}(X^1)} \cup V_{\alpha, \text{int}(X^2)} \cup W_\alpha$$

where $\text{int}(X^s)$ denotes the interior of the space X^s .

The portion V_{α, X^0} of M_α^3 consists of union of closed, connected components of M_α^3 which admit Riemannian metric of non-negative sectional curvature.

Proposition 3.4. *Let V_{α, X^0} be a union of one of following:*

- (1) *a spherical 3-dimensional space form;*
- (2) *a manifold double covered by $S^2 \times S^1$;*
- (3) *a closed flat 3-manifold.*

Then V_{α, X^0} can be collapsed to a 0-dimensional manifold with non-negative curvature. Moreover, V_{α, X^0} is a graph-manifold.

We denote the regular part of X^2 by X_{reg}^2 . Let us now recall a reduction for the proof of Perelman's collapsing theorem due to Morgan-Tian. Earlier related work on 3-dimensional collapsing theory was done by Xiaochun Rong in his thesis, (cf. [Rong93]).

Theorem 3.5 (Morgan-Tian [MT08], Compare [Rong93]). *Let $\{M_\alpha^3\}$ be a sequence of compact 3-manifolds satisfying the hypothesis of Theorem 0.1' and V_{α,X^0} be as in Proposition 3.4 above. If, for sufficiently large α , there exist compact, co-dimension 0 submanifolds $V_{\alpha,X^1} \subset M_\alpha^3$ and $V_{\alpha,X_{reg}^2} \subset M_\alpha^3$ with $\partial M_\alpha^3 \subset V_{\alpha,X^1}$ satisfying six conditions listed below, then Theorem 0.1 holds, where six conditions are:*

- (1) *Each connected component of V_{α,X^1} is diffeomorphic to the following.*
 - (a) *a T^2 -bundle over S^1 or a union of two twisted I -bundle over the Klein bottle along their common boundary;*
 - (b) *$T^2 \times I$ or $S^2 \times I$, where $I = [a, b]$ is a closed interval;*
 - (c) *a compact 3-ball or the complement of an open 3-ball in $\mathbb{RP}^3$ which is homeomorphic to $\mathbb{RP}^2 \times [0, \frac{1}{2}]$;*
 - (d) *a twisted I -bundle over the Klein bottle, or a solid torus.**In particular, every boundary component of V_{α,X^1} is either a 2-sphere or a 2-torus.*
- (2) $V_{\alpha,X^1} \cap V_{\alpha,X_{reg}^2} = (\partial V_{\alpha,X^1}) \cap (\partial V_{\alpha,X_{reg}^2})$;
- (3) *If N_0^2 is a 2-torus component of $\partial V_{\alpha,X^1}$, then $N_0^2 \subset \partial V_{\alpha,X_{reg}^2}$ if and only if N_0^2 is not boundary of ∂M_α^3 ;*
- (4) *If N_0^2 is a 2-sphere component of $\partial V_{\alpha,X^1}$, then $N_0^2 \cap \partial V_{\alpha,X_{reg}^2}$ is diffeomorphic to an annulus;*
- (5) V_{α,X_{reg}^2} *is the total space of a locally trivial S^1 -bundle and the intersection $V_{\alpha,X^1} \cap V_{\alpha,X_{reg}^2}$ is saturated under this fibration;*
- (6) *The complement $W_\alpha = [M_\alpha^3 - (V_{\alpha,X^0} \cup V_{\alpha,X^1} \cup V_{\alpha,X_{reg}^2})]$ is a disjoint union of solid tori and solid cylinders. The boundary of each solid torus is a boundary component of V_{α,X_{reg}^2} , and each solid cylinder $D^2 \times I$ in W_α meets V_{α,X^1} exactly in $D^2 \times \partial I$.*

Proof. ([MT08]) The proof of Theorem 3.5 is purely topological, which has nothing to do with the collapsing theory.

Morgan and Tian [MT08] first verified Theorem 3.5 for special cases under additional assumption on V_{α,X^1} :

- (i) V_{α,X^1} has no closed components;
- (ii) Each 2-sphere component of $\partial V_{\alpha,X^1}$ bounds a 3-ball component of V_{α,X^1} ;
- (iii) Each 2-torus component of $\partial V_{\alpha,X^1}$ that is compressible in M_α^3 bounds a solid torus component of V_{α,X^1} .

The general case can be reduced to a special case by a purely topological argument. $\square$

Definition 3.6. *If a collapsed 3-manifold M_α^3 has a decomposition*

$$M_\alpha^3 = V_{\alpha, X^0} \cup V_{\alpha, X^1} \cup V_{\alpha, X_{reg}^2} \cup W_\alpha$$

satisfying six properties listed in Theorem 3.5 and if V_{α, X^0} is a union of closed 3-manifolds which admit smooth Riemannian metrics of non-negative sectional curvature, then such a decomposition is called an admissible decomposition of M_α^3 .

In Section 1-2, we already discussed the part $V_{\alpha, \text{int}(X^2)}$ and a portion of W_α . In next section, we discuss the collapsing part $V_{\alpha, \text{int}(X^1)}$ with spherical or toral fibers for our 3-manifold M_α^3 , where $\text{int}(X^s)$ is the interior of X^s .

4. COLLAPSING WITH SPHERICAL AND TORAL FIBERS

In this section, we discuss the case when a sequence of metric balls $\{(B_{(M_\alpha^3, \tilde{g}_{ij}^\alpha)}(x_\alpha, r), x_\alpha)\}$ collapse to 1-dimensional space (X^1, x_∞) . There are only two choices of X^1 , either diffeomorphic to a circle or an interval $[0, l]$.

By Perelman's fibration theorem [Per94] or Yamaguchi's fibration theorem, we can find an open neighborhood U_α of x_α such that, for sufficiently large α , there is a fibration

$$N_\alpha^2 \rightarrow U_\alpha \rightarrow \text{int}(X^1)$$

where $\text{int}(X^1)$ is isometric to a circle S^1 or an open interval $(0, l)$.

We will use the soul theory (e.g., Theorem 2.11) to verify that a finite cover of the collapsing fiber N_α^2 must be homeomorphic to either a 2-sphere S^2 or a 2-dimensional torus T^2 , (see Figure 3 in §0)

Let us begin with two examples of collapsing 3-manifold with toral fibers

Example 4.1. Let $M^3 = \mathbb{H}^3/\Gamma$ be an oriented and non-compact quotient of $\mathbb{H}^3$ such that M^3 has finite volume and M^3 has exactly one end. Suppose that $\sigma : [0, \infty) \rightarrow \mathbb{H}^3/\Gamma$ be a geodesic ray. We consider the corresponding Busemann function $h_\sigma(x) = \lim_{t \rightarrow +\infty} [t - d(x, \sigma(t))]$. For sufficiently large c , the sup-level subset $V_{c, X^1} = h_\sigma^{-1}([c, +\infty))$ has special properties. It is well known that, in this case, the cusp end V_{c, X^1} is diffeomorphic to $T^2 \times [c, +\infty)$. Of course, the component $V_{c, X^1} \cong T^2 \times [c, +\infty)$ admits a collapsing family of metric $\{g_\varepsilon\}$, such that $(V_{c, X^1}, g_\varepsilon)$ is convergent to half line $[c, +\infty)$. $\square$

We would like to point out that 2-dimensional collapsing fibers can be collapsed at two different speeds.

Example 4.2. Let $M_\varepsilon^3 = (\mathbb{R}/\varepsilon\mathbb{Z}) \times (\mathbb{R}/\varepsilon^2\mathbb{Z}) \times [0, 1]$ be the product of rectangle torus $T_{\varepsilon, \varepsilon^2}$ and an interval. Let us fix a parametrization of $M_1^3 = \{(e^{2\pi si}, e^{2\pi ti}, u) | u \in [0, 1], s, t \in \mathbb{R}\}$ and $g_\varepsilon = \varepsilon^2 ds^2 + \varepsilon^4 dt^2 + du^2$. Then the re-scaled pointed spaces $\{((M_1^3, \frac{1}{\varepsilon^2}g_\varepsilon), (1, 1, \frac{1}{2}))\}$ are convergent to the limiting space (Y_∞^2, y_∞) , where Y_∞^2 is isometric to $S^1 \times (-\infty, +\infty)$.

Similarly, when the collapsing fiber is homeomorphic to a 2-sphere S^2 , the collapsing speeds could be different along longitudes and latitudes. We may assume that latitudes shrink at speed ε^2 and longitudes shrink at speed ε . For the same reason, after re-scaling, the limit space Y_∞^2 could be isometric to $[0, 1] \times (-\infty, +\infty)$. Thus, the non-compact limiting space Y_∞^2 could have boundary. $\square$

Let us return to the proof of Perelman's collapsing theorem. According to the second condition of Theorem 0.1, we consider a boundary component $N_{\alpha, X^1}^2 \subset \partial M_\alpha^2$, where the diameter of N_{α, X^1}^2 is at most $\omega_\alpha \rightarrow 0$ as $\alpha \rightarrow \infty$. Moreover, there exists a topologically trivial collar $\hat{V}_{\alpha, X^1}$ of length one and sectional curvatures of M_α^3 are between $(-\frac{1}{4} - \varepsilon)$ and $(-\frac{1}{4} + \varepsilon)$. In this case, we have a trivial fibration:

$$T_\alpha^2 \rightarrow \hat{V}_{\alpha, X^1} \xrightarrow{\pi_\alpha} [0, 1] \quad (4.1)$$

such that the diameter of each fiber $\pi_\alpha^{-1}(t)$ is at most $[\frac{\varepsilon + \varepsilon^{-1}}{2} \omega_\alpha]$ by standard comparison theorem. As $\alpha \rightarrow +\infty$, the sequence $\{\hat{V}_{\alpha, X^1}\}$ converge to an interval $X^1 = [0, 1]$.

We are ready to work on the main result of this subsection.

Theorem 4.3. *Let $\{((M_\alpha^3, \rho_\alpha^{-2} g_{ij}^\alpha), x_\alpha)\}$ be as in Theorem 0.1'. Suppose that an 1-dimensional space X^1 is contained in the 1-dimensional limiting space and $x_\alpha \rightarrow x_\infty$ is an interior point of X^1 . Then, for sufficiently large α , there exists a sequence of subsets $\hat{V}_{\alpha, X^1} \subset M_\alpha^3$ such that $\hat{V}_{\alpha, X^1}$ is fibering over $\text{int}(X^1)$ with spherical or toral fibers.*

$$N_\alpha^2 \rightarrow \hat{V}_{\alpha, \text{int}(X^1)} \rightarrow \text{int}(X^1)$$

where N_α^2 is homeomorphic to a quotient of a 2-sphere S^2 or a 2-torus T^2 .

When M_α^3 is oriented, N_α^2 is either S^2 or T^2 .

Proof. As we pointed out above, since $\text{int}(X^1)$ is a 1-dimensional space, there exists a fibration

$$N_\alpha^2 \rightarrow \hat{V}_{\alpha, \text{int}(X^1)} \rightarrow \text{int}(X^1)$$

for sufficiently large α . It remains to verify that the fiber is homeomorphic to $S^2, \mathbb{RP}^2, T^2$ or Klein bottle $T^2/\mathbb{Z}_2$. For this purpose, we use soul theory for possibly singular space Y_∞^k with non-negative curvature.

By our discussion, $B_{\rho_\alpha^{-2}g^\alpha}(x_\alpha, 1)$ is homeomorphic to $N_\alpha^2 \times (0, l)$, where N_α^2 is a closed 2-dimensional manifold. Thus, the distance function $r_{x_\alpha}(x) = d_{\rho_\alpha^{-2}g^\alpha}(x_\alpha, x)$ has at least one critical point $y_\alpha \neq x_\alpha$ in $B_{\rho_\alpha^{-2}g^\alpha}(x_\alpha, 1)$, because $B_{g^\alpha}(x_\alpha, \rho_\alpha)$ is not contractible. Let λ_α be

$$\max\{d_{\rho_\alpha^{-2}g^\alpha}(x_\alpha, y_\alpha) | y_\alpha \neq x_\alpha \text{ is a critical point of } r_{x_\alpha} \text{ in } B_{\rho_\alpha^{-2}g^\alpha}(x_\alpha, 1)\}.$$

We claim that $0 < \lambda_\alpha < 1$ and $\lambda_\alpha \rightarrow 0$ as $\alpha \rightarrow +\infty$. To verify this assertion, we observe that the distance functions $\{r_{x_\alpha}\}$ are convergent to $r_{x_\infty} : X^1 \rightarrow \mathbb{R}$. By Perelman's convergent theorem, the trajectory of gradient semi-flow

$$\frac{d^+\varphi}{dt} = \frac{\nabla r_{x_\alpha}}{|\nabla r_{x_\alpha}|^2}$$

is convergent to the trajectory in the limit space X^1 :

$$\frac{d^+\varphi}{dt} = \nabla r_{x_\infty} \quad (4.2)$$

(see [Petr07] or [KPT09]).

Clearly, $r_{x_\infty} : X^1 \rightarrow \mathbb{R}$ has no critical value in $(0, \delta_\infty)$ for $\delta_\infty > 0$. Thus, for sufficiently large α , the distance function r_{x_α} has no critical value in $(\lambda_\alpha, \delta_\infty - \varepsilon_\alpha)$, where $\lambda_\alpha \rightarrow 0$ and $\varepsilon_\alpha \rightarrow 0$ as $\alpha \rightarrow +\infty$.

Let us now re-scale our metrics $\{\rho_\alpha^{-2}g_{ij}^\alpha\}$ by $\{\lambda_\alpha^{-2}\}$ again. Suppose $\tilde{g}_{ij}^\alpha = \frac{1}{\lambda_\alpha^2 \rho_\alpha^2} g_{ij}^\alpha$. Then a subsequence of the sequence of the pointed spaces $\{(M_\alpha^3, \tilde{g}_{ij}^\alpha), x_\alpha\}$ will converge to (Y_∞^k, y_∞) . The curvature of Y_∞^k is greater than or equal to 0, because $\text{curv}_{\frac{1}{\lambda_\alpha^2 \rho_\alpha^2} g_{ij}^\alpha} \geq -\lambda_\alpha^2 \rightarrow 0$ as $\alpha \rightarrow +\infty$.

Since the distance $r_{y_\infty}(y) = d(y, y_\infty)$ has a critical point $z_\infty \neq y_\infty$ with $d(z_\infty, y_\infty) \leq 1$. Thus $\dim(Y_\infty^k) > 1$.

When $\dim(Y_\infty^k) = 3$, we observe that Y_∞^3 has exactly two ends in our case. Thus Y_∞^3 admits a line and hence its metric splits. A soul N_∞^2 of Y_∞^3 must be of 2-dimensional. Thus Y_∞^3 is isometric to $N_\infty^2 \times \mathbb{R}$. It follows that the soul N_∞^2 of Y_∞^3 has non-negative curvature. By Perelman's stability theorem, N_α^2 is homeomorphic to N_∞^2 for sufficiently large α . A closed possibly singular surface N_∞^2 of non-negative curvature has been classified in Section 2: $S^2, \mathbb{RP}^2, T^2$ or Klein bottle $T^2/\mathbb{Z}_2$.

Let us consider the case of $\dim(Y_\infty^k) = 2$. We may assume that the limiting space Y_∞^2 has exactly two ends. When Y_∞^2 has no boundary, then the limit space is isometric to $S^1 \times \mathbb{R}$, because Y_∞^2 has exactly two ends. By our discussion in §1-2, we have fibration structure

$$S^1 \rightarrow M_\alpha^3 \rightarrow Y_\infty^2 \quad (4.3)$$

for sufficiently large α .

When Y_∞^2 has non-empty boundary (i.e., $\partial Y_\infty^2 \neq \emptyset$), there are two subcases. If $y_\infty \in \text{int}(Y_\infty^2)$, by our discussion in Section 2, we still have $N_\alpha^2 \sim T^2/\Gamma$. If $y_\infty \in \partial Y_\infty^2$, using Theorem 5.4 below, we see that $N_\alpha^2 \sim [\partial B_{M_\alpha^3}(x_\alpha, r)] \sim S^2$. This completes the proof of our theorem for all cases. $\square$

In next section, we will discuss the end points or ∂X^1 when $X^1 \cong [0, l]$ is an interval. In addition, we also discuss the 2-dimensional boundary ∂X^2 when X^2 is a surface with convex boundary.

5. COLLAPSING TO BOUNDARY POINTS OR ENDPOINTS OF THE LIMITING SPACE

Suppose that a sequence of 3-manifolds is collapsing to a lower dimensional space X^s . In previous sections we showed that there is (possibly singular) fibration

$$N_\alpha^{3-s} \rightarrow B_{M_\alpha^3}(x_\alpha, r) \xrightarrow{G-H} \text{int}(X^s).$$

In this section we will consider the points on the boundary of X^s , we will divided our discussion into two cases: namely (1) when $s = 1$ and X^1 is a closed interval; and (2) when $s = 2$ and X^2 is a surface with boundary.

5.1. Collapsing to a surface with boundary.

Since X^2 is a topological manifold with boundary, without loss of generality, we can assume that $\partial X^2 = S^1$. First we provide two examples to demonstrate that the collapsing could occur in many different ways.

Example 5.1. Let $M_\frac{1}{\varepsilon}^3 = D^2 \times S^1$ be a solid torus, where $D^2 = \{(x, y) | x^2 + y^2 < 1\}$. We will construct a family of metrics g_ε on the disk so that $D_\varepsilon^2 = (D^2, g_\varepsilon)$ is converging to the interval $[0, 1]$. It follows that if $M_\frac{1}{\varepsilon}^3 = D_\varepsilon^2 \times S^1$ then the sequence of 3-spaces is converging to a finite cylinder: i.e., $M_\frac{1}{\varepsilon}^3 \rightarrow ([0, 1] \times S^1)$ as $\varepsilon \rightarrow 0$.

For this purpose, we let

$$D_\varepsilon^2 = \{(x, y, z) \in \mathbb{R}^3 | z^2 = \tan(\frac{1}{\varepsilon})[x^2 + y^2], z \geq 0, x^2 + y^2 + z^2 < 1\}$$

We further make a smooth perturbation around the vertex $(0, 0, 0)$, while keeping curvature non-negative. Clearly, as $\varepsilon \rightarrow 0$, the family of conic surfaces $\{D_\varepsilon^2\}$ collapses to an interval $[0, 1]$. Let $X^2 = [0, 1] \times S^1$ be the limiting space. As $\varepsilon \rightarrow 0$, our 3-manifolds $M_\frac{1}{\varepsilon}^3 = D_\varepsilon^2 \times S^1$ collapsed to X^2 with $\partial X^2 \neq \emptyset$.

In this example, for every point $p \in \partial X^2$, the space of direction $N_p(X^2)$ is a closed half circle with diameter π .

Example 5.2. In this example, we will construct the limiting surface with boundary corner points. Let us consider the unit circle $S^1 = \mathbb{R}/\mathbb{Z}$ and let $\varphi : \mathbb{R}^1 \rightarrow \mathbb{R}^1$ be an involution given by $\varphi(s) = -s$. Then its quotient $S^1/\langle\varphi\rangle$ is isometric to $[0, \frac{1}{2}]$. Our target limiting surface will be $X^2 = [0, 1) \times [0, \frac{1}{2}]$. Clearly, X^2 has boundary corner point with total angle $\frac{\pi}{2}$.

To see that X^2 is a limit of smooth Riemannian 3-manifolds $\{\bar{M}_{\frac{1}{\varepsilon}}^3\}$ while keeping curvatures non-negative, we proceed as follows. We first viewed a 2-sheet cover M^3 as an orbit space of N^4 by a circle action. Let $N^4 = D^2 \times (\mathbb{R}^2/(\mathbb{Z} \oplus \mathbb{Z}))$ and let $\{(re^{i\theta}, s, t) | 0 \leq \theta \leq 2\pi, 0 \leq s \leq 1, 0 \leq t \leq 1\}$ be a parametrization of $D^2 \times \mathbb{R}^2$ and hence for its quotient N^4 . There is a circle action $\psi_\lambda : N^4 \rightarrow N^4$ given by $\psi_\lambda(re^{i\theta}, s, t) = (re^{i(\theta+2\pi\lambda)}, s, t + \lambda)$ for each $e^{i2\pi\lambda} \in S^1$. We also define an involution $\tau : N^4 \rightarrow N^4$ by $\tau(re^{i\theta}, s, t) = (re^{i\theta}, -s, t + \frac{1}{2})$. It follows that $\tau \circ \tau = id$. Let $S^1 \ltimes \mathbb{Z}_2$ be subgroup generated by $\{\tau, \psi_\lambda\}_{\lambda \in S^1}$. We introduce a family of metrics:

$$g_\varepsilon = dr^2 + rd\theta^2 + ds^2 + \varepsilon^2 dt^2.$$

The transformations $\{\tau, \psi_\lambda\}_{\lambda \in S^1}$ remain isometries for Riemannian manifolds (N^4, g_ε) . Thus, $\bar{M}_{\frac{1}{\varepsilon}}^3 = (N^4, g_\varepsilon)/(S^1 \ltimes \mathbb{Z}_2)$ is a smooth Riemannian 3-manifold with non-negative curvature. It is easy to see that $\bar{M}_{\frac{1}{\varepsilon}}^3$ is homeomorphic to a solid torus. As $\varepsilon \rightarrow 0$, our Riemannian manifolds $\{\bar{M}_{\frac{1}{\varepsilon}}^3\}$ collapse to a lower dimensional space $X^2 = [0, 1) \times [0, \frac{1}{2}] = \{(r, s) | 0 \leq r < 1, 0 \leq s \leq \frac{1}{2}\}$. The surface X^2 has a corner point with total angle $\frac{\pi}{2}$.

In above two examples, if we set $\alpha = \frac{1}{\varepsilon}$, then there exist an open subset $U_\alpha \subset M_\alpha^3$ and a continue map $G_\alpha : U_\alpha \rightarrow X^2$ such that $G_\alpha^{-1}(A) = U_\alpha$ is homeomorphic to a solid torus, where A is an annular neighborhood of ∂X^2 in X^2 .

Let us recall an observation of Perelman on the distance function $r_{\partial X^2}$ from the boundary ∂X^2 .

Lemma 5.3. *Let X^2 be a compact Alexandrov surface of curvature $\geq c$ and with non-empty boundary ∂X^2 , $\Omega_{-\varepsilon} = \{p \in X^2 | d(p, \partial X^2) \geq \varepsilon\}$ and $A_\varepsilon = [X^2 - \Omega_{-\varepsilon}]$. Then, for sufficiently small ε , the distance function $r_{\partial X^2} = d_{X^2}(\cdot, \partial X^2)$ from the boundary has no critical point in A_ε .*

Proof. We will use a calculation of Perelman and a result of Petrunin to complete the proof. By the definition, if X^2 has curvature $\geq c$, then ∂X^2 must be convex. For any $x \in \partial X^2$, we let $\theta(x) = \text{diam}[\Sigma_x(X^2)]$ be the total tangent angle of the convex domain X^2 . It follows from the convexity $0 < \theta(x) \leq \pi$.

We consider the distance function $f(y) = d_{X^2}(y, \partial X^2)$ for all $y \in X^2$. Perelman (cf. [Per91] page 33, line 1) calculated that

$$|\nabla f(x)| = \sin\left(\frac{\theta(x)}{2}\right) > 0,$$

for all $x \in \partial X^2$. (Perelman stated his formula for spaces with non-negative curvature, but his proof using the first variational formula is applicable to our surface X^2 with curvature ≥ -1).

Corollary 1.3.5 of [Petr07] asserts that if there is a converging sequence $\{x_n\} \subset X^2$ with $x_n \rightarrow x \in \partial X^2$ as $n \rightarrow \infty$ then

$$\liminf_{n \rightarrow \infty} |\nabla f(x_n)| \geq |\nabla f(x)|.$$

Since X^2 is compact, it follows from the above discussion that there is an $\varepsilon > 0$ such that $|\nabla f(y)| > 0$ for $y \in A_\varepsilon$. $\square$

We now recall a theorem of Shioya-Yamaguchi with our own proof. The proof of Shioya-Yamaguchi used a version of Margulis Lemma, which we will use our Theorem 1.17 in §1 instead.

Theorem 5.4 ([SY00] page 2). *Let $\{(M_\alpha^3, p_\alpha)\}$ be a sequence of collapsing 3-manifolds as in Theorem 0.1'. Suppose that $\{(B_{M_\alpha^3}(p_\alpha, r), p_\alpha)\} \rightarrow (X^2, p_\infty)$ with $p_\infty \in \partial X^2$ and ∂X^2 is homeomorphic to S^1 . Then there is $\delta_1 > 0$ such that $B_{M_\alpha^3}(p_\alpha, \delta_1)$ is homeomorphic to $D^2 \times I \cong D^3$ for all sufficiently large α .*

Moreover, there exist an $\varepsilon > 0$ and a sequence of closed curves $\varphi_\alpha : S^1 \rightarrow M_\alpha^3$ with $\{\varphi_\alpha(S^1)\} \rightarrow \partial X^2$ as $B_{M_\alpha^3}(p_\alpha, r) \xrightarrow{G-H} X^2$ such that a ε -tubular neighborhood $U_\varepsilon[\varphi_\alpha(S^1)]$ of $\varphi_\alpha(S^1)$ in M_α^3 is homeomorphic to a solid tori $D^2 \times S^1$ for sufficiently large α .

Proof. By conic lemma (cf. Proposition 1.7), the number of points $p \in \partial X^2$ with $\text{diam}(\Sigma_p) \leq \frac{\pi}{2}$ is finite, denoted by $\{b_1, \dots, b_s\}$. Since $\partial X^2 \sim S^1$ is compact, it follows from Proposition 1.15 that there is a common $\ell > 0$ such that (1) the distance function $r_p(y) = d(p, y)$ has not critical point in $[B_{X^2}(p, \ell) - \{p\}]$; and (2) $B_{X^2}(p, \ell)$ is homeomorphic to the upper half disk D_+^2 , for all $p \in \partial X^2$.

Since ∂X^2 is homeomorphic to S^1 , we can approximate ∂X^2 by a sequence of closed broken geodesics $\{\sigma_\infty^{(j)}\}$ with vertices $\{q_1, \dots, q_j\} \subset \partial X^2$. We may assume that $\{b_1, \dots, b_s\}$ is a subset of $\{q_1, \dots, q_j\}$ for

all $j \geq s$. We also require that the distance between two consecutive vertices is less than c/j (i.e., $d_{\partial X^2}(q_i, q_{i+1}) \leq \frac{c}{j} < \frac{\ell}{8}$), for sufficiently large j .

We now choose a sequence of finite sets $\{p_1^\alpha, \dots, p_j^\alpha\}$ such that $p_i^\alpha \rightarrow q_i$ as $\alpha \rightarrow \infty$ and $\{p_1^\alpha, \dots, p_j^\alpha\}$ span an embedded broken geodesic $\sigma_\alpha^{(j)}$ in M_α^3 . It follows that

$$\sigma_\alpha^{(j)} \rightarrow \sigma_\infty^{(j)} \quad (5.1)$$

as $\alpha \rightarrow \infty$. Therefor, we may assume that there is a sequence of smooth embedded curves $\varphi_\alpha : S^1 \rightarrow M_\alpha^3$ such that

$$\varphi_\alpha(S^1) \rightarrow \partial X^2 \quad (5.2)$$

as $M_\alpha^3 \rightarrow X^2$. Thus, when $\{z(B_{M_\alpha^3}(p_\alpha, r), p_\alpha)\} \rightarrow (X^2, p_\infty)$, the corresponding closed curves $\varphi_\alpha(S^1) \xrightarrow{G-H} \partial X^2$ in the Gromov-Hausdorff topology, as $M_\alpha^3 \rightarrow X^2$.

We now choose a sufficiently large j_0 and divide our closed curve $\varphi_\alpha : S^1 \rightarrow M_\alpha^3$ into j_0 -many arcs of constant speed:

$$\varphi_{\alpha,i} : [a_i, a_{i+1}] \rightarrow M_\alpha^3$$

for $i = 0, 1, \dots, j_0$, where $\varphi_\alpha(a_{j_0+1}) = \varphi_\alpha(a_0)$.

For a fixed i , we let

$$A_{\alpha,s}^i = \varphi_\alpha([a_i + s, a_{i+1} - s])$$

and $q_{\alpha,i} = \varphi_\alpha(a_i)$. We will show that there is an ε_i such that

$$U_{\varepsilon_i}(A_{\alpha,0}^i) = B_{M_\alpha^3}(q_{\alpha,i}, \varepsilon_i) \cup U_{\varepsilon_i}(A_{\alpha,s}^i) \cup B_{M_\alpha^3}(q_{\alpha,i+1}, \varepsilon_i) \quad (5.3)$$

is homeomorphic to the unit 3-ball D^3 . Our theorem will follow from (5.3). It remains to establish (5.3).

We will show that each of $\{B_{M_\alpha^3}(q_{\alpha,i}, \varepsilon_i), U_{\varepsilon_i}(A_{\alpha,s}^i), B_{M_\alpha^3}(q_{\alpha,i+1}, \varepsilon_i)\}$ is homeomorphic to D^3 .

For $U_{\varepsilon_i}(A_{\alpha,s}^i)$, we first observe that ∂X^2 is convex. Thus, the boundary arc $\varphi_\infty([a_i + s, a_{i+1} - s]) \subset \partial X^2$ is a Perelman-Sharafutdinov semi-gradient curve of the distance function $r_{q_{\infty,i}}(y) = d_{X^2}(y, q_{\infty,i})$. We already choose ℓ sufficiently small and j_0 sufficiently large so that $r_{q_{\infty,i}}$ has no critical point on $[B_{X^2}(q_{\infty,i}, \ell) - \{q_{\infty,i}\}]$. By our construction, we have $r_{q_{\alpha,i}}(\cdot) \rightarrow r_{q_{\infty,i}}(\cdot)$, as $\alpha \rightarrow \infty$. It follows from Proposition 1.14 that the distance function has no critical points on $U_{\varepsilon_i}(A_{\alpha,s}^i)$. Using Perelman's fibration theorem, we obtain a fibration

$$N_{\alpha,i}^2 \rightarrow U_{\varepsilon_i}(A_{\alpha,s}^i) \xrightarrow{r_{q_{\alpha,i}}} (s, \ell_{\alpha,i} - s).$$

It follows that $U_{\varepsilon_i}(A_{\alpha,s}^i) \sim [N_{\alpha,i}^2 \times (s, \ell_{\alpha,i} - s)]$. We will first use Theorem 1.17 and its proof to show that $\partial N_{\alpha,i}^2 \sim S^1$.

Let ε_0 be given by Lemma 5.3 and $r_\alpha(y) = d_{M_\alpha^3}(y, \varphi_\alpha(S^1))$. By the proof of Theorem 1.17, the map $F_\alpha(z) = (r_{q_{\alpha,i}}(z), r_\alpha(z))$ is regular on $Z_{\alpha,i}^3(s/2, \varepsilon_0) = [U_{\varepsilon_0}(A_{\alpha,s}^i) - U_{s/2}(A_{\alpha,s}^i) - B_{M_\alpha^3}(q_{\alpha,i}, \varepsilon_i) - B_{M_\alpha^3}(q_{\alpha,i+1}, \varepsilon_i)]$. There is a circle fibration $S^1 \rightarrow Z_{\alpha,i}^3(\frac{s}{2}, \varepsilon_0) \xrightarrow{F_\alpha} \mathbb{R}$. This proves that $\partial N_{\alpha,i}^2 \sim S^1$. It follows that $H_{\alpha,i}^2 = \{\partial[U_{\varepsilon_i}(A_{\alpha,s}^i)] - B_{M_\alpha^3}(q_{\alpha,i}, \varepsilon_i) - B_{M_\alpha^3}(q_{\alpha,i+1}, \varepsilon_i)\}$ is homeomorphic to a cylinder $S^1 \times (s, \ell_{\alpha,i} - s)$. Using two points suspension of the cylinder $H_{\alpha,i}^2$, we can further show that

$$\partial[U_{\varepsilon_i}(A_{\alpha,s}^i)] \sim S^2$$

It remains to show that $U_{\varepsilon_i}(A_{\alpha,s}^i) \sim D^3$ for sufficiently small $\varepsilon_i > 0$. Suppose contrary, we argue as follows. Using Proposition 1.14, we see that r_α has no critical points in $[U_{\varepsilon_0}(A_{\alpha,s}^i) - U_{\varepsilon_0/2}(A_{\alpha,s}^i)]$. Let λ_α be the largest critical value of r_α in $U_{\varepsilon_0}(A_{\alpha,s}^i)$. By our assumption, $\lambda_\alpha \rightarrow 0$ as $\alpha \rightarrow \infty$. We now consider a sequence of re-scaled spaces $\{(\frac{1}{\lambda_\alpha} M_\alpha^3, q_{\alpha,i})\}$. Its sub-sequence converges to a limiting space $(Y_\infty^s, \bar{q}_{\infty,i})$. Recall that $\dim(X^2) = 2$. For the same reason as in the proof of Proposition 2.3, we can show that $\dim(Y_\infty^s) = 3$ and that Y_∞^3 has no boundary. Let $\bar{N}_\infty^s$ be the soul of Y_∞^3 . There are three possibilities.

(1) If the soul $\bar{N}_\infty^s$ is a point, then by Theorem 2.11 we obtain that $Y_\infty^3 \sim \mathbb{R}^3$. It follows that $U_{\varepsilon_i}(A_{\alpha,s}^i) \sim D^3$ by Perelman's stability theorem, we are done.

(2) If the soul $\bar{N}_\infty^s$ is a circle, then Y_∞^3 (or its double cover) is isometric to $S^1 \times Z^2$, where Z^2 is homeomorphic to $\mathbb{R}^2$. Let $\bar{\varphi}_\infty(\mathbb{R})$ be the limit curve in the re-scaled limit space Y_∞^3 . Since Y_∞^3 (or its double cover) is isometric to $S^1 \times Z^2$, we have $U_r(\bar{\varphi}_\infty([-R, R]))$ is homeomorphic to $S^1 \times D^2$. By Perelman's stability theorem, we would have $\partial[U_{\varepsilon_i}(A_{\alpha,s}^i)] \sim \partial[S^1 \times D^2] \sim T^2$, which contradicts to the assertion $\partial[U_{\varepsilon_i}(A_{\alpha,s}^i)] \sim S^2$.

(3) If the soul $\bar{N}_\infty^s$ of Y_∞^3 has dimension 2, then it follows from that the infinity of Y_∞^3 would have at most two points. However, since $\dim(X^2) = 2$, the infinity of Y_∞^3 has an arc, a contradiction. This completes the proof of $U_{\varepsilon_i}(A_{\alpha,s}^i) \sim D^3$ for sufficiently large α .

With extra efforts, we can also show that $B_{M_\alpha^3}(q_{\alpha,i}, \varepsilon) \sim D^3$ for sufficiently large α . Hence, $U_\varepsilon[\varphi_\alpha(S^1)] \sim \cup_i D_i^3 \sim [D^2 \times S^1]$. This completes our proof. $\square$

5.2. Collapsing to a closed interval.

Since all our discussions in this sub-section are semi-local, we may have the following setup:

$$\lim_{\alpha \rightarrow +\infty} (M_\alpha^3, p_\alpha) = (I, O) \quad (5.4)$$

in the pointed Gromov-Hausdorff distance, and $I = [0, \ell]$ is an interval, $O \in I$ is an endpoint of I . We will study the topology of $B_{M_\alpha^3}(p_\alpha, r)$ for a given small r .

We begin with four examples to illustrate how smooth Riemannian 3-manifolds M_α^3 collapse to an interval $[0, \ell]$ with curvature bounded from below. Collapsing manifolds in these manifolds are homeomorphic to one of the following: $\{D^3, [\mathbb{RP}^3 - D^3], S^1 \times D^2, K^2 \times [0, \frac{1}{2}]\}$, where $D^3 = \{x \in \mathbb{R}^3 \mid |x| < 1\}$ and K^2 is the Klein bottle.

Example 5.5. (M_α^3 is homeomorphic to D^3). For each $\varepsilon > 0$, we consider a convex hypersurface in $\mathbb{R}^4$ as follows. We glue a lower half of the 3-sphere

$$S_{\varepsilon,-}^3 = \{(x_1, x_2, x_3, x_4) \in \mathbb{R}^4 \mid 0 \leq x_4 \leq \varepsilon, x_1^2 + x_2^2 + x_3^2 + (x_4 - \varepsilon)^2 = \varepsilon^2\}$$

to a finite cylinder

$$S_\varepsilon^2 \times [\varepsilon, 1] = \{(x_1, x_2, x_3, x_4) \in \mathbb{R}^4 \mid \varepsilon \leq x_4 \leq 1, x_1^2 + x_2^2 + x_3^2 = \varepsilon^2\}.$$

Our 3-manifold $M_\varepsilon^3 = S_{\varepsilon,-}^3 \cup (S_\varepsilon^2 \times [\varepsilon, 1])$ collapse to the unit interval as $\varepsilon \rightarrow 0$.

For other cases, we consider the following example.

Example 5.6. (M_α^3 homeomorphic to $S^1 \times D^2$). Let us glue a lower half of 2-sphere

$$S_{\varepsilon,-}^2 = \{(x_1, x_2, x_3) \in \mathbb{R}^3 \mid 0 \leq x_3 \leq \varepsilon, x_1^2 + x_2^2 + (x_3 - \varepsilon)^2 = \varepsilon^2\}$$

to a finite cylinder $S_\varepsilon^1 \times [\varepsilon, 1]$. The resulting disk

$$D_\varepsilon^2 = S_{\varepsilon,-}^2 \cup (S_\varepsilon^1 \times [\varepsilon, 1])$$

is converge to unit interval, as $\varepsilon \rightarrow 0$. We could choose $M_{\frac{1}{\varepsilon}}^3 = S_{\varepsilon^2,-}^1 \times D_\varepsilon^2$.

It is clear that $M_{\frac{1}{\varepsilon}}^3 \rightarrow [0, 1]$ as $\varepsilon \rightarrow 0$.

We now would like to consider the remaining cases. Of course, two un-oriented surfaces $\mathbb{RP}_\varepsilon^2 = S_\varepsilon^2/\mathbb{Z}^2$ and $K^2 = T^2/\mathbb{Z}_2$ would converge to a point, as $\varepsilon \rightarrow 0$. However, the *twisted* I -bundle over $\mathbb{RP}^2$ (or K^2) is homeomorphic to an oriented manifold $\mathbb{RP}^2 \times [0, \frac{1}{2}] = [\mathbb{RP}^3 - D^3]$ (or $K^2 \times [0, \frac{1}{2}] = M\ddot{o} \times S^1$), where $M\ddot{o}$ is the Möbius band.

Example 5.7. (M_α^3 homeomorphic to $\mathbb{RP}^2 \times [0, \frac{1}{2}]$ or $M\ddot{o} \times S^1$).

Let us first consider round sphere $S_\varepsilon^2 = \{(x_1, x_2, x_3) \in \mathbb{R}^3 \mid |x| = \varepsilon\}$. There is an orientation preserving involution $\tau : \mathbb{R}^3 \times [-\frac{1}{2}, \frac{1}{2}]$ given by $\tau(x, t) = (-x, -t)$. Suppose that $\langle \tau \rangle = \mathbb{Z}_2$ is the subgroup generated by τ . Thus, the quotient of $S_\varepsilon^2 \times [-\frac{1}{2}, \frac{1}{2}]$ is an orientable manifold $\mathbb{RP}^2 \times [0, \frac{1}{2}] = [\mathbb{RP}^3 - D^3]$.

Similarly, we can consider the case of $K_\varepsilon^2 \times [0, \frac{1}{2}] \rightarrow [0, \frac{1}{2}]$, where K_ε^2 is a Klein bottle.

Shioya and Yamaguchi showed that the above examples exhausted all cases up to homeomorphisms.

Theorem 5.8 (Theorem 0.5 of [SY00]). *Suppose that $\lim_{\alpha \rightarrow +\infty} (M_\alpha^3, p_\alpha) = (I, O)$ with curvature ≥ -1 and $I = [0, \ell]$. Then M_α^3 is homeomorphic to a gluing of C_0 and C_ℓ to $N^2 \times (0, 1)$, where C_0 and C_ℓ are homeomorphic to one of $\{D^3, [\mathbb{RP}^3 - D^3], S^1 \times D^2, K^2 \times [0, \frac{1}{2}]\}$ and N^2 is a quotient of T^2 or S^2 .*

For the proof of Theorem 5.8, we need to establish two preliminary results (see Theorem 5.9 - 5.10 below). Let us consider possible exceptional orbits in the Seifert fibration

$$S^1 \rightarrow M_\alpha^3 \rightarrow \text{int}(Y^2) \quad (5.5)$$

for sufficiently large α . We emphasize that the topological structure of M_α^3 depends on the number of extremal points (or called essential singularities) of Y^2 in this case. Moreover, the topological structure of M_α^3 also depends on the type of essential singularity of Y^2 , when we glue a pair of solid tori together, (see (5.16) below). Therefore, we need the following theorem with a new proof.

Theorem 5.9 (Compare with Corollary 14.4 of [SY00]). *Let Y^2 be a connected, non-compact and complete surface with non-negative curvature and with possible boundary. Then the following is true.*

- (i) *If Y^2 has no boundary, then Y^2 has at most two extremal points (or essential singularities). When Y^2 has exactly two extremal points, Y^2 is isometric to the double $\text{dbl}([0, \ell] \times [0, \infty))$ of the flat half strip.*
- (ii) *If Y^2 has non-empty boundary $\partial Y^2 \neq \emptyset$, then Y^2 has at most one interior essential singularity.*

Proof. For (i), we will use the multi-step Perelman-Sharafutdinov semi-flows to carry out the proof. For the assertion (ii), we consider the Cheeger-Gromoll type Busemann function

$$f(y) = \lim_{t \rightarrow \infty} [t - d(y, \partial B_{Y^2}(y_0, t))]. \quad (5.6)$$

In Section 2, we already showed that $\Omega_c = f^{-1}((-\infty, c])$ is compact for any finite c . Let $a_0 = \inf\{f(y) | y \in Y^2\}$. Then the level set $\Omega_{a_0} = F^{-1}(a_0)$ has dimension at most 1. Recall that Ω_{a_0} is convex by the soul theory. Thus, Ω_{a_0} is either a point or isometric to a length-minimizing geodesic segment: $\sigma_0 : [0, \ell] \rightarrow Y^2$.

Case a. If $\Omega_{a_0} = \{z_0\}$ is a point soul of Y^2 , by an observation of Grove [Grv93] we see that the distance function $r(y) = d(y, z_0)$ has no critical point in $[Y^2 - \{z_0\}]$. It follows that z_0 is only possible extremal point of Y^2 .

Case b. If $\Omega_{a_0} = \sigma_0([0, \ell])$, then for $s \in (0, \ell)$ Petrunin [Petr07] showed that $T_{\sigma_0(s)}^-(Y^2) = \mathbb{R}^2$. Hence, two endpoints $\sigma_0(0)$ and $\sigma_0(\ell)$ are only two possible extremal points of Y^2 .

Suppose that Y^2 has exactly two extremal points $\sigma_0(0)$ and $\sigma_0(\ell)$. We choose two geodesic rays $\psi_0 : [0, \infty) \rightarrow Y^2$ and $\psi_\ell : [0, \infty) \rightarrow Y^2$ from two extremal points $\sigma_0(0)$ and $\sigma_0(\ell)$ respectively. The broken geodesic $\Gamma = \sigma_0 \cup \psi_0 \cup \psi_\ell$ divides our space Y^2 into two connected components:

$$[Y^2 - \Gamma] = \Omega_- \cup \Omega_+. \quad (5.7)$$

We now consider the distance function

$$w_\pm(u) = d_{\Omega_\pm}(\psi_\ell(u), \psi_0(u)). \quad (5.8)$$

Let us consider the Perelman-Sharafutdinov semi-gradient flow $\frac{d^+w}{dt} = \nabla(-f)|_w$ for Busemann function $-f(w)$. Recall that the semi-flow is distance non-increasing, since the curvature is non-negative. Hence, we see that $t \rightarrow w_\pm(u - t)$ is a non-increasing function of $t \in [0, u]$. It follows that

$$w_\pm(u) \geq w_\pm(0) = \ell \quad (5.9)$$

for $u \geq 0$. On other hand, we could use multi-step geodesic triangle comparison theorem as in [CDM09] to verify that

$$w_\pm(u) \leq w_\pm(0) = \ell \quad (5.10)$$

with equality holds if and only if four points $\{\sigma_0(0), \sigma_0(\ell), \psi_\ell(u), \psi_0(u)\}$ span a flat rectangle in $\Omega_\pm$. Therefore, $\Omega_\pm$ is isometric to $[0, \ell] \times [0, \infty)$. It follows that Y^2 is isometric to the double $\text{dbl}([0, \ell] \times [0, \infty))$.

The second assertion (ii) follows from (i) by using $\text{dbl}(Y^2)$. $\square$

For compact surfaces X^2 of non-negative curvature, we use an observation of Perelman ([Per91] page 31) together with multi-step Perelman-Sharafutdinov semi-flows, in order to estimate the number of extremal points in X^2 .

Theorem 5.10. (1) *Suppose that X^2 is a compact and oriented surface with non-negative curvature and with non-empty boundary $\partial X^2 \neq \emptyset$. Then X^2 has at most two interior extremal points. When X^2 has two interior extremal points, then X^2 is isometric to a gluing of two copies of flat rectangle along their three corresponding sides.*

(2) *Suppose that X^2 is a closed and oriented surface with non-negative curvature. Then X^2 has at most four extremal points.*

Proof. (1) We consider the double $\text{dbl}(X^2)$ of X^2 . If $\text{dbl}(X^2)$ is not simply-connected and if it is an oriented surface with non-negative curvature, then we already showed that $\text{dbl}(X^2)$ is a flat torus.

We may assume that X^2 is homeomorphic to a disk: $X^2 \cong D^2$ and $\partial X^2 = S^1$. Let $X_{-\varepsilon} = \{x \in X^2 \mid d(x, \partial X^2) \geq \varepsilon\}$. Perelman [Per91] already showed that $X_{-\varepsilon}$ remains to be convex, (see also [CDM09]). If $a_0 = \max\{d(x, \partial X^2)\}$, then X_{-a_0} is either a geodesic segment or a single point set. Thus, X^2 has at most two interior extremal points. The rest of the proof is the same as that of Theorem 5.9 with minor modifications.

(2) Suppose that X^2 has 2 distinct extremal points p and q . Let N be a geodesic segment connecting p and q . We need to show for the function $f(x) = d_N(x) = d(N, x)$ is concave for all $x \in X^2 \setminus N$. Clearly for $x \in X^2 \setminus N$ there exists $x_N \in N$ such that $|xx_N| = d(x, N)$. There are two possibilities:

- (i) x_N is in the interior of N , then proof of the concavity of f is exactly same as the one of Theorem 6.1 in [Per91] (see also [Petr07] p156 and [CDM09]).
- (ii) x_N is one of the endpoints, say p , then by first variational formula we have

$$d_{\Sigma_p}(\uparrow_p^x, \uparrow_p^q) \geq \frac{\pi}{2} \quad (5.11)$$

where $\uparrow_p^x$ denotes the set of directions of geodesics from p to x in Σ_p . On the other hand, by our assumption p is an interior extremal point so

$$\text{diam} \Sigma_p \leq \frac{\pi}{2} \quad (5.12)$$

Combine (5.11) and (5.12) we have


$$d_{\Sigma_p}(\uparrow_p^x, \uparrow_p^q) = \frac{\pi}{2}$$

The rest of the proof is same as the one of Theorem 6.1 in [Per91] (or [Petr07], [CDM09]).

Hence, we have shown that f is concave function on $X^2 \setminus N$. Let A be the maximum set. Then Ω_1 is either a geodesic segment or one point. Thus, $X^2 \setminus N$ contains at most 2 extremal points by our proof of (1). Therefore, the number of total extremal points is at most 4. $\square$

The case of exactly 4 extremal points on a topological 2-sphere can be illustrated in Figure 8. Let us now complete the proof of Theorem 5.8.

Proof of Theorem 5.8:

FIGURE 8. A 2-sphere S^2 with 4 essential singularities

The proof is due to Shioya-Yamaguchi [SY00]. Our new contribution is the simplified proof of Theorem 5.9, which will be used in the study of subcases (1.b) and (2.b) below. For the convenience of readers, we provide a detailed argument here.

Recall that in Section 4, we already constructed a fibration

$$N_\alpha^2 \rightarrow U_\alpha \rightarrow \text{int}(X^1)$$

with shrinking fiber N_α^2 is homeomorphic to either S^2 or T^2 for sufficiently large α .

When $\{p_\alpha\} \rightarrow p_\infty \in \partial X^1$, since the fibers $\{N_\alpha^2\}$ are shrinking, we may assume that

$$\partial B_{M_\alpha^3}(p_\alpha, \delta) = S^2 \quad \text{or} \quad \partial B_{M_\alpha^3}(p_\alpha, \delta) = T^2 \quad (5.13)$$

for sufficiently large α , where $X^1 = [0, \ell]$ and $\delta = \ell/2$.

If there exists an ε_0 such that $r_\alpha(x) = d_{M_\alpha^3}(x, p_\alpha)$ has no critical points on the punctured ball $[B_{M_\alpha^3}(p_\alpha, \varepsilon_0) - \{p_\alpha\}]$ for sufficiently large α , then, using the proof of Proposition 1.7, we can show that $B_{M_\alpha^3}(p_\alpha, \varepsilon_0) \sim D^3$ for sufficiently large α . Thus, conclusion of Theorem 5.8 holds for this case.

Otherwise, there exists a subsequence $\{\varepsilon_\alpha\} \rightarrow 0$ such that $r_\alpha(x) = d_{M_\alpha^3}(x, p_\alpha)$ has a critical point q_α with $d(p_\alpha, q_\alpha) = \varepsilon_\alpha$. It follows that $\left(\frac{1}{\varepsilon_\alpha} M_\alpha^3, \bar{p}_\alpha\right) \rightarrow (Y^k, \bar{p}_\infty)$. Using a similar argument as in the proof of Proposition 2.3, we can show that the limiting space Y^k has non-negative curvature and $\dim(Y^k) = k \geq 2$, where $\bar{p}_\alpha$ is the image of p_α under the re-scaling. There are two cases described in (5.13) above.

Case 1. $\partial B_{M_\alpha^3}(p_\alpha, \delta) = S^2$.

If $B_{M_\alpha^3}(p_\alpha, \delta)$ is homeomorphic to D^3 , then no further verification is needed. Otherwise, we may assume $B_{M_\alpha^3}(p_\alpha, \delta) \neq D^3$ and $\partial B_{M_\alpha^3}(p_\alpha, \delta) = S^2$.

Under these two assumptions, we would like to verify that $B_{M_\alpha^3}(p_\alpha, \delta)$ is homeomorphic to $\mathbb{RP}^2 \times I$. There are two sub-cases:

Subcase 1.a. If $\dim Y = 3$, then Y^3 is non-negatively curved, open and complete. Let N^k be a soul of Y^3 .

Using Perelman's stability theorem, we claim that the soul N^k cannot be S^1 . Otherwise the boundary of $B_{M_\alpha^3}(p_\alpha, \delta)$ would be T^2 , a contradiction. For the same reason, the soul N^k cannot be T^2 or K^2 . Otherwise the boundary $\partial B_{M_\alpha^3}(p_\alpha, \delta)$ would be homeomorphic to T^2 , contradicts to our assumption. Because the boundary of $B_{M_\alpha^3}(p_\alpha, \delta)$ only consists of one component, the soul of N^k of Y cannot be S^2 ; otherwise we would have $Y^3 \sim S^2 \times \mathbb{R}$ and $\partial B_{M_\alpha^3}(p_\alpha, \delta) \sim S^2 \cup S^2$, a contradiction.

Therefore, we have demonstrated that the soul N^k must be either a point or $\mathbb{RP}^2$. It follows that either $B_Y(N^k, R) \cong D^3$ or $\mathbb{RP}^2 \times I$ by soul theorem. Hence, we conclude that $B_Y(N^k, R) \cong \mathbb{RP}^2 \times I$ for sufficiently large R . Using Perelman's stability theorem, we conclude that $B_{M_\alpha^3}(p_\alpha, \delta)$ is homeomorphic to $\mathbb{RP}^2 \times I$ for this sub-case.

Subcase 1.b. If $\dim Y = 2$, i.e. Y is a surface with possibly non-empty boundary. First we claim $\partial Y \neq \emptyset$. For any fixed $r > 0$, we have

$$\partial B_{M_\alpha^3}(p_\alpha, r) \cong S^2 \quad (5.14)$$

by our assumption that the regular fiber is S^2 . Suppose contrary, $\partial Y^2 = \emptyset$. We would have $Y \cong \mathbb{R}^2$ (or a Möbius band) by Theorem 2.6, because Y^2 has one end. Thus, for sufficiently large R we have $\partial B_Y(\bar{p}_\infty, R) = S^1$. Applying fibration theorem for the collapsing to the surface case, we would further have

$$\partial B_{M_\alpha^3}(\bar{p}_\alpha, R) = T^2$$

which contradicts to our boundary condition (5.14). Hence, Y has non-empty boundary: $\partial Y^2 \neq \emptyset$.

Since Y^2 has one end and $\partial Y^2 \neq \emptyset$, by Corollary 2.9 we know that Y^2 is homeomorphic to either upper-half plane $[0, \infty) \times \mathbb{R}$ or isometric a half cylinder. By previous argument, Y^2 can not be isometric to a half cylinder. Hence, we have

$$Y^2 \cong [0, \infty) \times \mathbb{R} \quad (5.15)$$

and that ∂Y is a non-compact set.

We further observe that if $\bar{p}_\infty$ is a boundary point of Y^2 , then, by Theorem 5.4, we would have

$$B_{M_\alpha^3}(p_\alpha, r) \cong [D_+^2 \times I] \cong D^3$$

where D_+^2 is closed half disk, a contradiction.

Thus we may assume that $\bar{p}_\infty$ is an interior point of Y^2 . Let us consider the Seifert fiber projection $B_{\frac{1}{\lambda_\alpha}M_\alpha^3}(\bar{p}_\alpha, R) \rightarrow B_Y(\bar{p}_\infty, R)$ for some large R . If $B_Y(\bar{p}_\infty, R)$ contains no interior extremal point, then by the proof of Theorem 5.4 one would have $B_{\frac{1}{\varepsilon_\alpha}M_\alpha^3}(\bar{p}_\alpha, R) \cong D^2 \times I \cong D^3$, a contradiction. Therefore, there exists an extremal point inside $B(\bar{p}_\infty, R)$. Without loss of generality, we may assume this extremal point is O .

By Theorem 5.9 and the fact that $\partial Y^2 \neq \emptyset$, we observe that Y^2 has at most one interior extremal point. In our case, Y^2 is isometric to the following flat surface with a singularity. Let $\Omega = [0, h] \times [0, \infty)$ be a half flat strip and $\Gamma = \{(x, y) \in \Omega \mid xy = 0\} \subset \partial\Omega$. Our singular flat surface Y^2 is isometric to a gluing two copies of Ω along the curve Γ .

The picture of $B_Y(O, R)$ for large R will look like Figure 9, where the bold line denotes $\partial Y \cap B_Y(O, R)$. By our assumption, we have




FIGURE 9. A metric disk $B_{Y^2}(O, R)$ in Y^2 for large R

$\partial B(\bar{p}_\alpha, R) \cong S^2$. Thus we can glue a 3-ball D^3 to $B(\bar{p}_\alpha, R)$ along $\partial B(\bar{p}_\alpha, R) \cong S^2$ to get a new closed 3-manifold $\hat{M}_\alpha^3$.

Recall that we have a (possibly singular) fibration

$$S^1 \rightarrow B(\bar{p}_\alpha, R) \xrightarrow{G_\alpha} \text{int}(Y^2).$$

By Theorem 2.1, we have $G_\alpha^{-1}(B_{Y^2}(O, h/2)) \cong D^2 \times S^1$. With some efforts, we can further show that $[\hat{M}_\alpha^3 - G_\alpha^{-1}(B_{Y^2}(O, h/4))] \cong S^1 \times D^2$.

The exceptional orbit $G_\alpha^{-1}(O)$ is corresponding to the case of $m_0 = 2$ in Example 2.0. Finally we conclude that $\hat{M}_\alpha^3$ is homeomorphic to real projective 3-space:

$$\hat{M}_\alpha^3 \cong (D^2 \times S^1) \cup_{\psi_{\mathbb{Z}_2}} (S^1 \times D^2) \cong \mathbb{RP}^3. \quad (5.16)$$

Therefore, by our construction, we have

$$B_{M_\alpha^3}(p_\alpha, \delta) = [\hat{M}_\alpha^3 \setminus D^3] \cong [\mathbb{RP}^3 \setminus D^3] \cong \mathbb{RP}^2 \ltimes I.$$

This completes the first part of our proof for the case of $\partial B_{M_\alpha^3}(p_\alpha, \delta) = S^2$.

Case 2. If $\partial B_{M_\alpha^3}(p_\alpha, \delta) = T^2$, our discussion will be similar to the previous case with some modifications. In fact, there are still two sub-cases:

Subcase 2.a. If $\dim Y = 3$, then the Y has a soul N^k . We assert that N^k cannot be a point nor S^2 nor $\mathbb{RP}^2$ because the boundary $\partial B_{M_\alpha^3}(p_\alpha, \delta) = T^2$. In addition, we observe that N^k cannot be T^2 since the boundary of $B_{M_\alpha^3}(p_\alpha, \delta)$ consists only one component. Therefore, there are two remaining cases: N^k is homeomorphic to either S^1 or Klein bottle K^2 . If the soul is S^1 then $B_{M_\alpha^3}(p_\alpha, \delta) \cong D^2 \times S^1$. Similarly, if soul is K^2 , then $B_{M_\alpha^3}(p_\alpha, \delta) \cong K^2 \ltimes I$.

Subcase 2.b. $\dim Y = 2$. It is clear that our limiting space Y^2 is non-compact.

If $\partial Y^2 = \emptyset$, we proceed as follows. (i) When the soul of Y^2 is S^1 , by the connectedness of $\partial B_{M_\alpha^3}(p_\alpha, \delta)$ we know that Y is a open Möbius band. Therefore, $B_{M_\alpha^3}(p_\alpha, \delta)$ is homeomorphic to product of Möbius band and S^1 , i.e. a twist I -bundle over K^2 . (ii) When the soul of Y^2 is a single point, $Y = \mathbb{R}^2$, which is non-compact. By Theorem 5.9, we see that the number k of extremal points in Y^2 is at most 2. Recall that there is (possibly singular) fibration: $S^1 \rightarrow B_{M_\alpha^3}(p_\alpha, \delta) \xrightarrow{G_\alpha} \text{int}(Y^2)$. If $k \leq 1$, we have $B_{M_\alpha^3}(p_\alpha, \delta) \cong S^1 \times D^2$. If $k = 2$, by Theorem 5.9, we can further show that $B_{M_\alpha^3}(p_\alpha, \delta) \cong K^2 \ltimes I$.

Let us now handle the remaining subcase when $\partial Y^2 \neq \emptyset$ is not empty. By a similar argument as in subcase 1.b above, we conclude that $B_{M_\alpha^3}(p_\alpha, \delta)$ is homeomorphic to either $\mathbb{RP}^2 \ltimes I$ or D^3 , which contradicts to our assumption $\partial B_{M_\alpha^3}(p_\alpha, \delta) \cong T^2$.

This completes the proof of Theorem 5.8 for all cases. $\square$

6. GLUING LOCAL FIBRATION STRUCTURES AND CHEEGER-GROMOV'S COMPATIBILITY CONDITION

In this section, we complete the proof of Perelman's collapsing theorem for 3-manifolds (Theorem 0.1'). In previous five sections, we made progress in decomposing each collapsing 3-manifold M_α^3 into several parts:

$$M_\alpha^3 = V_{\alpha, X^0} \cup V_{\alpha, X^1} \cup V_{\alpha, \text{int}(X^2)} \cup W_\alpha \quad (6.1)$$

where V_{α, X^0} is a union of closed smooth 3-manifolds of non-negative sectional curvature, V_{α, X^1} is a union of fibrations over 1-dimensional spaces with spherical or toral fibers and $V_{\alpha, \text{int}(X^2)}$ admits locally defined almost free circle actions.

Extra cares are needed to specify the definition of V_{α, X^i} for each α . For example, we need to choose specific parameters for collapsing

3-manifolds $\{M_\alpha^3\}$ so that the decomposition in (6.1) becomes well-defined. The choices of parameters can be made in a similar way as in Section 3 of [SY05].

Theorem 6.0. *Suppose that $\{(M_\alpha^3, g^\alpha)\}$ be as in Theorem 0.1 and that M_α^3 has no connected components which admit metrics of non-negative curvature. Then there are two small constants $\{c_1, \varepsilon_1\}$ such that*

$$V_{\alpha, X^1} = \{x \in M_\alpha^3 \mid d_{GH}(B_{(M_\alpha^3, \rho_\alpha^{-2} g_{ij}^\alpha)}(x, 1), [0, \ell]) \leq \varepsilon_1\}$$

and

$$V_{\alpha, X^2} = \{x \in M_\alpha^3 \mid \text{Vol}(B_{g^\alpha}(x, \rho_\alpha(x))) \leq c_1 \varepsilon_1 [\rho_\alpha(x)]^3\}$$

which are described as in (6.1) above and $\{\rho_\alpha(x)\}$ satisfy inequality (6.8) below.

Proof. Recall that if the pointed Riemannian manifolds $\{((M_\alpha^3, \rho_\alpha^{-2} g_{ij}^\alpha), x_\alpha)\}$ converge to (X^k, x_∞) , then by our assumption we have $\text{diam}(X^k) \geq 1$ and hence

$$1 \leq \dim[X^k] \leq 2.$$

Therefore, there are two cases: (1) $\dim[X^k] = 1$ and (2) $\dim[X^k] = 2$.

Case 1. $\dim[X^k] = 1$. By the proof of Theorem 5.8, there exists $\varepsilon_1 > 0$ such that if

$$d_{GH}(B_{(M_\alpha^3, \rho_\alpha^{-2} g_{ij}^\alpha)}(x, 1), [0, \ell]) \leq \varepsilon_1 \quad (6.2)$$

then $B_{(M_\alpha^3, \rho_\alpha^{-2} g_{ij}^\alpha)}(x, 1)$ admits a possibly singular fibration over $[0, 1]$, where the curvature of the $\rho_\alpha^{-2} g_{ij}^\alpha$ is bounded below by -1 . Clearly, ℓ must satisfy $1 - \varepsilon_1 \leq \ell \leq 2 + \varepsilon_1$.

If the inequality (6.2) holds, then the unit metric ball is very thin and can be covered by at most $\frac{4\ell}{\varepsilon_1}$ many small metric balls $\{B_{(M_\alpha^3, \rho_\alpha^{-2} g_{ij}^\alpha)}(y_j, 2\varepsilon_1)\}$. By Bishop volume comparison theorem, we have a volume estimate:

$$\text{Vol}[B_{(M_\alpha^3, \rho_\alpha^{-2} g_{ij}^\alpha)}(x, 1)] \leq \frac{4\ell}{\varepsilon_1} c_0 [\sinh(2\varepsilon_1)]^3 \leq c_0^* \ell \varepsilon_1^2 \quad (6.3)$$

In this case, we set $x \in V_{\alpha, X^1}$.

Case 2. $\dim[X^k] = 2$. There are two subcases for

$$d_{GH}(B_{(M_\alpha^3, \rho_\alpha^{-2} g_{ij}^\alpha)}(x, 1), B_{X^2}(x_\infty, 1)) \leq \varepsilon_2 \quad (6.4)$$

Subcase 2a. The metric disk $B_{X^2}(x_\infty, 1)$ has small area.

In this subcase, one can choose constant $\varepsilon_2 > 0$ such that if a metric disk in X^2 with radius r satisfies $1/2 \leq r \leq 1$ and area $\leq \varepsilon_2$ then by comparison theorems one can prove

$$d_{GH}(B_{X^2}(x_\infty, 1), [0, \ell]) \leq \varepsilon_1/3 \quad (6.5)$$

for some $\ell > 0$. It follows that

$$\begin{aligned} d_{GH}(B_{(M_\alpha^3, \rho_\alpha^{-2} g_{ij}^\alpha)}(x_\alpha, 1), [0, \ell]) &\leq d_{GH}(B_{(M_\alpha^3, \rho_\alpha^{-2} g_{ij}^\alpha)}(x_\alpha, 1), B_{X^2}(x_\infty, 1)) \\ &\quad + d_{GH}(B_{X^2}(x_\infty, 1), [0, \ell]) \\ &\leq \frac{\varepsilon_1}{3} + \frac{\varepsilon_1}{3} < \varepsilon_1 \end{aligned}$$

We can now view the ball $B_{\rho_\alpha^{-2} g_{ij}^\alpha}(x_\alpha, 1)$ is fibred over $[0, \ell]$, instead of fibred over the disk $B_{X^2}(x_\infty, 1)$.

In this subcase, we let x_α be in both V_{α, X^1} and V_{α, X^2} .

Subcase 2b. The metric disk $B_{X^2}(x_\infty, 1)$ is very fat (thick).

In this subcase, we may assume that the length of metric circle $\partial B_{X^2}(x_\infty, r)$ is 100 times greater than ε_1 for $r \in [\frac{1}{2}, 1]$. Hence, the length of collapsing fiber $F_\alpha^{-1}(z)$ for $z \in \partial B_{X^2}(x_\infty, r)$. Thus, the metric spheres $\{\partial B_{(M_\alpha^3, \rho_\alpha^{-2} g_{ij}^\alpha)}(x_\alpha, r)\}$ collapse in only one direction. Because G_α is an almost metric submersion due to Perelman's semi-flow convergence theorem, (cf. Proposition 1.14 above), volumes of metric balls collapse at an order $o(\varepsilon_1)$:

$$\text{Vol}[B_{(M_\alpha^3, \rho_\alpha^{-2} g_{ij}^\alpha)}(x, r)] \leq c_1 \varepsilon_1 r^3 \quad (6.6)$$

for $r \in [\frac{1}{2}, 1]$. Moreover, the free homotopy class of collapsing fibers is *unique* in the annular region $A_{(M_\alpha^3, \rho_\alpha^{-2} g_{ij}^\alpha)}(x_\alpha, \frac{1}{4}, 1)$.

By the discussion above, we have the following decomposition of the manifold M_α^3 for α large. More precisely, for sufficiently large α , M_α^3 has decomposition

$$M_\alpha^3 = V_{\alpha, X^1} \cup V_{\alpha, \text{int}(X^2)} = V_{\alpha, \text{int}(X^1)} \cup V_{\alpha, \text{int}(X^2)} \cup W_\alpha$$

where W_α contains collapsing parts near ∂X^1 and ∂X^2 described in Section 5 above. This completes the proof. $\square$

Let us now recall an observation of Morgan-Tian about the choices of functions $\{\rho_\alpha(x)\}$ and volume collapsing factors $\{w_\alpha\}$.

Proposition 6.1 (Morgan-Tian [MT08]). *Let M_α^3, w_α and $\rho_\alpha(x)$ be as in Theorem 0.1'. Suppose that none of connected components of M_α^3 admits a smooth Riemannian metric of non-negative sectional curvature. Then, by changing w_α by a factor $\hat{c}$ independent of α , we can choose ρ_α so that*

$$\rho_\alpha(x) \leq \dim(M_\alpha^3) \quad (6.7)$$

and

$$\rho_\alpha(y) \leq \rho_\alpha(x) \leq 2\rho_\alpha(y) \quad (6.8)$$

for all $y \in B_{g_\alpha}(x, \frac{\rho_\alpha}{2})$

Proof. (cf. Proof of Lemma 1.5 of [MT08]) Since there is a typo in Morgan-Tian's argument, for convenience of readers, we reproduce their argument with minor corrections.

Let us choose

$$C_0 = \max_{0 \leq s \leq 1} \left\{ \frac{s^3}{4\pi \int_0^s (\sinh u)^2 du} \right\}$$

For each connected component $N_{\alpha,j}^3$ of M_α^3 , the Riemannian sectional curvature of $N_{\alpha,j}^3$ can not be everywhere non-negative. Thus, for each $x \in N_{\alpha,j}^3$, there is a maximum $r = r_\alpha(x) \geq \rho_\alpha(x)$ such that sectional curvature of g_α on $B_{g_\alpha}(x, r)$ is greater than or equal to $-\frac{1}{r^2}$. Consequently, curvature of $\frac{1}{r^2}g_\alpha$ on $B_{\frac{1}{r^2}g_\alpha}(x, 1)$ is ≥ -1 . By Bishop-Gromov relative comparison theorem and our assumption $\text{Vol}(B_{g_\alpha}(x, \rho_\alpha)) \leq w_\alpha \rho_\alpha^3$, we have

$$\begin{aligned} \text{Vol}(B_{g_\alpha}(x, r)) &= r^3 \text{Vol}[B_{\frac{1}{r^2}g_\alpha}(x, 1)] \leq r^3 \frac{V_{\text{hyp}}(1)}{V_{\text{hyp}}(\frac{\rho_\alpha}{r})} \text{Vol}(B_{\frac{1}{r^2}g_\alpha}(x, \frac{\rho_\alpha}{r})) \\ &= \frac{V_{\text{hyp}}(1)}{V_{\text{hyp}}(\frac{\rho_\alpha}{r})} \text{Vol}(B_{g_\alpha}(x, \rho_\alpha)) \leq \frac{V_{\text{hyp}}(1)}{V_{\text{hyp}}(\frac{\rho_\alpha}{r})} w_\alpha \rho_\alpha^3 \\ &= \frac{V_{\text{hyp}}(1)}{V_{\text{hyp}}(\frac{\rho_\alpha}{r})} \left(\frac{\rho_\alpha}{r}\right)^3 \frac{1}{(\frac{\rho_\alpha}{r})^3} w_\alpha \rho_\alpha^3 \leq C_0 V_{\text{hyp}}(1) w_\alpha r^3 \end{aligned} \tag{6.9}$$

where $V_{\text{hyp}}(s)$ is the volume of the ball $B_{\mathbb{H}^3}(p_0, s)$ of radius s in 3-dimensional hyperbolic space with constant negative curvature -1 .

Let $\hat{C} = C_0 V_{\text{hyp}}$ be a constant number independent of α , and Rm_g be the sectional curvature of the metric g . We now replace $\rho_\alpha(x)$ by

$$\rho_\alpha^*(x) = \max\{r \mid |Rm_{g_\alpha}|_{B_{g_\alpha}(x, r)} \geq -\frac{1}{r^2}\} \tag{6.10}$$

Our new choice $\rho_\alpha^*(x)$ clearly satisfies

$$\begin{cases} \rho_\alpha^*(x) \leq \text{diam}(M_\alpha^3) \\ \frac{1}{2}\rho_\alpha^*(x) \leq \rho_\alpha^*(y) \leq 2\rho_\alpha^*(x) \end{cases} \tag{6.11}$$

for all $y \in B_{g_\alpha}(x, \frac{\rho_\alpha^*}{2})$ and $x \in M_\alpha^3$. $\square$

Our next goal in this section is to show that we can perturb our decomposition above along their boundaries so that the new decomposition admits an F-structure in the sense of Cheeger-Gromov, and hence M_α^3 is a graph manifold for sufficiently large α . Let us begin with a special case.

6.1. Perelman's collapsing theorem for a special case.

In this subsection, we prove Theorem 0.1' for a special case when Perelman's fibrations are assumed to be circle fibrations or toral fibrations. In next sub-section, we reduce the general case to the special case, (i.e. the case when no spherical fibration occurred).

As we pointed out earlier, for the proof of Theorem 0.1 it is sufficient to verify that M_α^3 admits an F-structure of positive rank in the sense of Cheeger-Gromov for sufficiently large α , (cf. [CG86], [CG90], [Rong93]).

Recall that an F-structure on a 3-manifold M^3 is a collection of charts $\{(U_i, V_i, T^{k_i})\}$ such that T^{k_i} acts on a finite normal cover V_i of U_i and the T^{k_i} -action on V_i commutes with deck transformation on V_i . Moreover the tori-group actions satisfy a compatibility condition on any possible overlaps.

Let us state compatibility condition as follows.

Definition 6.2 (Cheeger-Gromov's compatibility condition).

Let $\{(U_i, V_i, T^{k_i})\}$ be a collection of charts as above. If, for any two charts (U_i, V_i, T^{k_i}) and (U_j, V_j, T^{k_j}) with non-empty intersection $U_i \cap U_j \neq \emptyset$ and with $k_i \leq k_j$, the T^{k_i} actions commutes with the T^{k_j} -actions on a finite normal cover of $U_i \cap U_j$ after re-parametrization if needed, then the collection $\{(U_i, V_i, T^{k_i})\}$ is said to satisfy Cheeger-Gromov's compatibility condition.

For the purpose of this paper, since our manifolds under consideration are 3-dimension, the choice of free tori T^{k_i} actions must be either circle action or 2-dimensional torus action. Thus we only have to consider following three possibilities:

- (i) Both overlapping charts (U_i, V_i, S^1) and (U_j, V_j, S^1) admit almost free circle actions;
- (ii) Both overlapping charts (U_i, V_i, T^2) and (U_j, V_j, T^2) admit almost free torus actions;
- (iii) There are a circle action (U_i, V_i, S^1) and a torus-action (U_j, V_j, T^2) with non-empty intersection $U_i \cap U_j \neq \emptyset$.

Let us begin with the sub-case (ii).

Proposition 6.3. *Let U_{α, i_1} and U_{α, i_2} be two overlapping open subsets in U_{α, X^1} with toral fibers $F_{i_1}^2 \cong F_{i_2}^2 \cong T^2$ described in Section 4. Suppose that $U_{\alpha, i_1} \cap U_{\alpha, i_2} \neq \emptyset$. Then we can modify charts $(U_{\alpha, i_1}, V_{\alpha, i_1}, T^2)$ so that the perturbed torus-actions on modified charts satisfy the Cheeger-Gromov's compatibility condition.*

Proof. Recall that in Section 4 we worked on the following diagrams

$$\begin{array}{ccc}
 U_{\alpha,i_1} & \xrightarrow{f_{i_1}} & \mathbb{R} \\
 \downarrow \text{G-H} & & \uparrow \\
 X_{i_1}^1 & \xrightarrow{r\rho_{i_1}} & \mathbb{R}
 \end{array}
 \quad \text{and} \quad
 \begin{array}{ccc}
 U_{\alpha,i_2} & \xrightarrow{f_{i_2}} & \mathbb{R} \\
 \downarrow \text{G-H} & & \uparrow \\
 X_{i_2}^1 & \xrightarrow{r\rho_{i_2}} & \mathbb{R}
 \end{array}$$

Without loss of generality, we may assume that $X_{i_1} = (a_{i_1}, b_{i_1})$ and $X_{i_2} = (a_{i_2}, b_{i_2})$ with non-empty intersection $X_{i_1} \cap X_{i_2} = (a_{i_2}, b_{i_1}) \neq \emptyset$, where $a_{i_1} < a_{i_2} < b_{i_1} < b_{i_2}$. Let $\{\lambda_1, \lambda_2\}$ be a partition of unity of $[a_{i_1}, b_{i_2}]$ corresponding to the open cover $\{(a_{i_1}, b_{i_1}), (a_{i_2}, b_{i_2})\}$. After choosing $\pm$ sign and λ_1, λ_2 carefully, we may assume that

$$\hat{f}(t) = \lambda_1(t)f_{i_1}(t) \pm \lambda_2(t)f_{i_2}(t)$$

will not have any critical point $t \in [a_{i_1}, b_{i_2}]$. We also require that $\hat{f} : X_{i_1} \cup X_{i_2} \rightarrow \mathbb{R}$ is a *regular* function in the sense of Perelman (i.e. $\hat{f}$ is a composition of distance function given by Definition 1.8-1.9). Thus, we can lift the admissible function $\hat{f}$ to a function

$$f_\alpha : U_{\alpha,i_1} \cup U_{\alpha,i_2} \rightarrow \mathbb{R}$$

such that

$$\nabla f_\alpha|_y \neq 0$$

for $y \in [U_{\alpha,i_1} \cup U_{\alpha,i_2}]$, $f_\alpha \cong f_{\alpha,i_1}$ on $[U_{\alpha,i_1}^* - U_{\alpha,i_2}^*]$ and $f_\alpha \cong f_{\alpha,i_2}$ on $[U_{\alpha,i_2}^* - U_{\alpha,i_1}^*]$, where U_{α,i_1}^* is a perturbation of U_{α,i_1} and U_{α,i_2}^* is a perturbation of U_{α,i_2} . Thus there is a new perturbed torus fibration.

$$T^2 \rightarrow U_{\alpha,i_1} \cup U_{\alpha,i_2} \rightarrow Y^1.$$

This completes the proof. $\square$

Similarly, we can modify admissible maps to glue two circle actions together.

Proposition 6.4. *Let U_{α,i_1} and U_{α,i_2} be two open sets contained in U_{α,X_{reg}^2} corresponding to a decomposition of M_α^3 described in Section 1. Suppose that two charts $\{(U_{\alpha,i_1}, V_{\alpha,i_1}, S^1), (U_{\alpha,i_2}, V_{\alpha,i_2}, S^1)\}$ have non-empty overlap $U_{\alpha,i_1} \cap U_{\alpha,i_2} \neq \emptyset$. Then the union $U_{\alpha,i_1} \cup U_{\alpha,i_2}$ admits a global circle action after some modifications when needed.*

Proof. We may assume that, in the limiting processes of $U_{\alpha,i_1} \rightarrow X_1^2$ and $U_{\alpha,i_2} \rightarrow X_2^2$, both limiting surfaces X_1^2 and X_2^2 are fat (having relatively large area growth). For the remaining cases when either X_1^2 and X_2^2 are relatively thin, by the proof of Theorem 6.0 we can view either

U_{α,i_1} or U_{α,i_1} is a portion of V_{α,X^1} instead. These remaining situations can be handled by either Proposition 6.3 above or Proposition 6.5 below respectively.

As we pointed out in Section 1-2 the exceptional orbits are isolated. Without loss of generality, we may assume that there is no exceptional circle orbits occurs in the overlap $U_{\alpha,i_1} \cap U_{\alpha,i_2}$.

Since both limiting surfaces X_1^2 and X_2^2 are very *fat*, the lengths of metric circles $\partial B_{X_i^2}(x_{\infty,i}, r)$ is at least 100 times larger than ε_1 , which is great than the length of collapsing fibers, where $r \in [\frac{1}{2}, 1]$. Hence, on the overlapping region $U_{\alpha,i_1} \cap U_{\alpha,i_2}$, two collapsing fibres are freely homotopic each other in the shell-type region $A_{(M_\alpha^3, \rho_\alpha^{-2} g_{ij}^\alpha)}(x_\alpha, \frac{1}{4}, 1)$.

Let us consider the corresponding diagrams again.

$$\begin{array}{ccc} U_{\alpha,i_1} & \xrightarrow{\psi_{i_1}} & \mathbb{R}^2 \\ \downarrow \text{G-H} & & \uparrow \\ X_{i_1}^2 & \longrightarrow & \mathbb{R}^2 \end{array} \quad \text{and} \quad \begin{array}{ccc} U_{\alpha,i_2} & \xrightarrow{\psi_{i_2}} & \mathbb{R}^2 \\ \downarrow \text{G-H} & & \uparrow \\ X_{i_2}^2 & \longrightarrow & \mathbb{R}^2 \end{array}$$

We can use 2×2 matrix-valued function $A(x)$ to construct a new *regular* map

$$\psi_\alpha : U_{\alpha,i_1} \cup U_{\alpha,i_2} \rightarrow \mathbb{R}^2$$

from $\{\psi_{\alpha,i_1}, \psi_{\alpha,i_2}\}$ as follows. Let

$$\psi_\alpha = \lambda_1 A_1 \psi_{i_1} + \lambda_2 A_2 \psi_{i_2}$$

where $\{\lambda_1, \lambda_2\}$ is a partition of unity for U_{i_1}, U_{i_2} , $A_{i_1}(x)$ and $A_{i_2}(x)$ are 2×2 matrix-valued functions such that

- (i) $A_{i_1}(x) \cong \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ is close to the identity on $[U_{i_1}^* - U_{i_2}^*]$.
- (ii) Similarly, $A_{i_2}(x) \cong \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ is close to the identity on $[U_{i_2}^* - U_{i_1}^*]$.

where $U_{i_1}^*$ is a perturbation of U_{i_1} and $U_{i_2}^*$ is a perturbation of U_{i_2} . We leave the details to readers. $\square$

We now discuss relations between circle actions and torus action.

Proposition 6.5. *Let $U_{\alpha,i_1} \subset U_{\alpha,X^1}$ and $U_{\alpha,i_2} \subset U_{\alpha,X_{reg}^2}$, where $\{U_{\alpha,X^1}, U_{\alpha,X_{reg}^2}, U_{\alpha,X^0}, W_\alpha\}$ is a decomposition of M_α^3 as in Section 1-5. Suppose that $(U_{\alpha,i_1}, V_{\alpha,i_1}, T^2)$ and $(U_{\alpha,i_2}, V_{\alpha,i_2}, S^1)$ have an interface $U_{\alpha,i_1} \cap U_{\alpha,i_2} \neq \emptyset$. Then, after a perturbation if need, the circle orbits are contained in torus orbits on the overlap.*

Proof. Let $f_{\alpha,i_1} : U_{\alpha,i_1} \rightarrow \mathbb{R}$ be the regular function which induces the chart $(U_{\alpha,i_1}, V_{\alpha,i_1}, T^2)$. Suppose that $\psi_{\alpha,i_1} = (f_{\alpha,i_2}, g_{\alpha,i_2}) : U_{\alpha,i_2} \rightarrow \mathbb{R}^2$ is the corresponding regular map. Since any component of a regular map must be regular, after necessary modifications, we may assume that $f_{\alpha,i_1}^* : U_{\alpha,i_1}^* \rightarrow \mathbb{R}$ is equal to f_{α,i_2}^* in the modified regular map $\psi_{\alpha,i_2}^* = (f_{\alpha,i_2}^*, g_{\alpha,i_2}^*) : U_{\alpha,i_2}^* \rightarrow \mathbb{R}^2$ on the overlap $U_{\alpha,i_1}^* \cap U_{\alpha,i_2}^*$. It follows that the modified S^1 -orbits are contained in the newly perturbed T^2 -orbits. Thus, the modified charts $(U_{\alpha,i_1}^*, V_{\alpha,i_1}^*, T^2)$ and $(U_{\alpha,i_2}^*, V_{\alpha,i_2}^*, S^1)$ satisfy the Cheeger-Gromov's compatibility condition. $\square$

We now conclude this sub-section by a special case of Perelman's collapsing theorem.

Theorem 6.6. *Suppose that $\{(M_\alpha^3, g_\alpha)\}$ satisfies all conditions stated in Theorem 0.1. Suppose that all Perelman fibrations are either (possible singular) circle fibrations or toral fibrations. Then M_α^3 must admits a F-structure of positive rank in the sense of Cheeger-Gromov for sufficiently large α . Consequently, M_α^3 is a graph manifold for sufficiently large α .*

Proof. By our assumption and Proposition 6.1, there is a collection of charts $\{(U_{\alpha,i}, V_{\alpha,i}, T^{k_i})\}_{i=1}^n$ such that

- (i) $1 \leq k_i \leq 2$;
- (ii) $\{U_{\alpha,i}\}_{i=1}^n$ is an open cover of M_α^3 ;
- (iii) $(U_{\alpha,i}, V_{\alpha,i}, T^{k_i})$ satisfies condition (3.1).

It remains to verify that our collection

$$\{(U_{\alpha,i}, V_{\alpha,i}, T^{k_i})\}_{i=1}^n$$

satisfies the Cheeger-Gromov compatibility condition, after some modifications. Since exceptional circle orbits with non-zero Euler number are isolated, we may assume that on any possible overlap

$$U_{\alpha,i_1} \cap \cdots \cap U_{\alpha,i_j} \neq \emptyset$$

there is no exceptional circular orbits. Applying Proposition 6.3-6.5 when needed, we can perturb our charts so that the modified collection of charts $\{(U_{\alpha,i}^*, V_{\alpha,i}^*, T^{k_i})\}_{i=1}^n$ satisfy the Cheeger-Gromov compatibility condition. It follows that M_α^3 admits an F-structure $\mathcal{F}^*$ of positive rank. Therefore, M_α^3 is a graph manifold for sufficiently large α . $\square$

6.2. Perelman's collapsing theorem for general case.

Our main difficulty is to handle a Perelman fibration with possible spherical fibers:

$$S^2 \rightarrow U_{\alpha,i} \rightarrow \text{int}(X^1)$$

because the Euler number of S^2 is non-zero and S^2 does not admit any free circle actions.

Proposition 6.7. *Let $\{(M_\alpha^3, g_\alpha)\}$ be a sequence of Riemannian 3-manifolds as in Theorem 0.1. If there is a Perelman fibration*

$$N^2 \rightarrow U_{\alpha, X^1} \rightarrow (a, b)$$

with spherical fiber $N^2 \cong S^2$, then U_{α, X^1} must be contained in the interior of M_α^3 when M_α^3 has non-empty boundary $\partial M_\alpha^3 \neq \emptyset$.

Proof. According to condition (2) of Theorem 0.1, for each boundary component $N_{\alpha, j}^2 \subset \partial M_\alpha^3$, there is a topologically trivial collar $V_{\alpha, j}$ of length one such that $V_{\alpha, j}$ is diffeomorphic to $T^2 \times [0, 1]$. Thus, we have $U_{\alpha, j} \cap \partial M_\alpha^3 = \emptyset$ $\square$

We need to make the following elementary but useful observation.

Proposition 6.8. (1) *Let $A = \{(x_1, x_2, x_3) \in \mathbb{R}^3 \mid |\vec{x}| = 1, |x_3| \leq \frac{1}{2}\}$ be an annulus. The product space $S^2 \times [0, 1]$ has a decomposition*

$$(D_+^2 \times [0, 1]) \cup (A \times [0, 1]) \cup (D_-^2 \times [0, 1]),$$

where $D_\pm^2 = \{(x_1, x_2, x_3) \in S^2(1) \mid \pm x_3 \geq \frac{1}{2}\}$.

(2) *If $\{(M_\alpha^3, g_\alpha)\}$ satisfies conditions of Theorem 0.1, then, for sufficiently large α , M_α^3 has a decomposition $\{U_{\alpha, j}\}_{j=1}^{m_\alpha}$ such that*

(2.a) *either a finite normal cover $V_{\alpha, j}$ of $U_{\alpha, j}$ admits a free T^{k_i} -action with $k_i = 1, 2$.*

(2.b) *or $U_{\alpha, j}$ is homeomorphic to a finite solid cylinder $D^2 \times [0, 1]$ with $U_{\alpha, j} \cap \partial M_\alpha^3 = \emptyset$.*

(3) *If $U_{\alpha, j}$ is a finite cylinder as in (2.b), then it must be contained in a chain $\{U_{\alpha, j_1}, \dots, U_{\alpha, j_m}\}$ of finite solid cylinders such that their union*

$$\hat{W}_{\alpha, h_j} = \bigcup_{i=1}^m U_{\alpha, j_i}$$

is homeomorphic to a solid torus $D^2 \times S^1$.

Proof. The first two assertions are trivial. It remains to verify the third assertion. By our construction, if $U_{\alpha, j}$ is a finite cylinder homeomorphic to $D^2 \times I$, then $U_{\alpha, j}$ meets V_{α, X^1} exactly in $D^2 \times \partial I$. Moreover, such a finite solid cylinder $U_{\alpha, j}$ is contained in a chain $\{U_{\alpha, j_1}, \dots, U_{\alpha, j_m}\}$ of solid cylinders. It is easy to see that the union $\hat{W}_{\alpha, h_j}$ is homotopic to its core S^1 . i.e. $\hat{W}_{\alpha, h_j}$ is homeomorphic to a solid torus $D^2 \times S^1$. $\square$

We are ready to complete the proof of Perelman's collapsing theorem.

Proof of Theorem 0.1. Using Proposition 6.1 we can choose ρ_α so that

$$\rho_\alpha(x) \leq \dim(M_\alpha^3)$$

and

$$\rho_\alpha(y) \leq \rho_\alpha(x) \leq 2\rho_\alpha(y)$$

for all $y \in B_{g_\alpha}(x, \frac{\rho_\alpha}{2})$. Therefore, it is sufficient to verify Theorem 0.1' instead.

By our discussion above, for sufficiently large α , our 3-manifold M_α^3 admits a collection of (possibly singular) Perelman fibration. Thus M_α^3 has a decomposition

$$M_\alpha^3 = V_{\alpha, X^0} \cup V_{\alpha, X^1} \cup V_{\alpha, \text{int}(X^2)} \cup W_\alpha$$

For each chart in $V_{\alpha, X^0} \cup V_{\alpha, \text{int}(X^2)}$, it admits a Seifert fibration. However, remaining charts could be homeomorphic to $S^2 \times I$, $[\mathbb{RP}^3 - D^3]$ or a solid cylinder $D^2 \times I$. It follows from Proposition 6.8(3) that M_α^3 has a more refined decomposition

$$M_\alpha^3 = \bigcup_{i=1}^m U_{\alpha, j}$$

such that

- (i) either a finite normal cover $V_{\alpha, j}$ of $U_{\alpha, j}$ admits a free T^{k_j} -action with $k_j = 1, 2$;
- (ii) or $U_{\alpha, j}$ is homeomorphic to a solid torus $D^2 \times S^1$, which is obtained by a chain of solid cylinders.

Observe that possibly exceptional circular orbits are isolated. Moreover if $\{U_{\alpha, j}\}$ are of type (2.b) in Proposition 6.8, these cores $\{0\} \times S^1$ are isolated as well.

It remains to verify that our collection of charts $\{(U_{\alpha, j}, V_{\alpha, j}, T^{k_j})\}$ satisfy Cheeger-Gromov compatibility condition. By observations on exceptional orbits and cores of solid cylinders, we may assume that on possibly overlap

$$U_{\alpha, j_1} \cap \cdots \cap U_{\alpha, j_k} \neq \emptyset$$

there is no exceptional orbits nor cores of solid cylinders.

We require that $V_{\alpha, \text{int}(X^2)}$ meets S^2 -factors as annular type A_α . Thus, if $U_{\alpha, j} \cong S^1 \times D^2$, we can introduce T^2 -actions on $\partial U_{\alpha, j} \cong S^1 \times S^1$ which are compatible with S^1 -actions on A_α . After modifying our charts as in proofs of Proposition 6.3-6.5, we can obtain a new collection of charts $\{(U_{\alpha, j}^*, V_{\alpha, j}^*, T^{k_j})\}$ satisfying the Cheeger-Gromov's compatibility condition. Therefore, M_α^3 admits an F-structure $\mathcal{F}_\alpha$ of positive rank. It follows that M_α^3 is a 3-dimensional graph manifold, (cf. [Rong93]). $\square$

Acknowledgement: Both authors are grateful to Professor Karsten Grove for teaching us the modern version of critical point theory for distance functions. Professor Takashi Shioya carefully proofread every sub-step of our simple proof in the entire paper, pointing out several inaccurate statements in an earlier version. He also generously provided us a corrected proof of Proposition 2.2. We are very much indebted to Professor Xiaochun Rong for supplying Theorem 6.0 and its proof. Hao Fang and Christina Sormani made useful comments on an earlier version. Finally, we also appreciate Professor Brian Smyth's generous help the exposition in our paper. We thank the referee for his (or her) suggestions, which led many improvements.

REFERENCES

- [BBBMP10] L. Bessières, G. Besson, M. Boileau, S. Maillot, J. Porti, *Collapsing irreducible 3-manifolds with nontrivial fundamental group* Invent. Math. **179** (2010) 434-460.
- [BBI01] Dima Burago, Yuri Burago; S. Ivanov, *A course in metric geometry*. Graduate Studies in Mathematics, **33**. American Mathematical Society, Providence, RI, 2001. xiv+415 pp. ISBN: 0-8218-2129-6.
- [BGP92] Yu Burago; M. Gromov; G. Perelman, *A. D. Aleksandrov spaces with curvatures bounded below*. (Russian) Uspekhi Mat. Nauk **47** (1992), no. 2(284), 3–51, 222; translation in Russian Math. Surveys **47** (1992), no. 2, 1–58
- [CalC92] E. Calabi and J. Cao, *Simple closed geodesics on convex surfaces*. Journal of Differential Geometry **36** (1992), 517-549.
- [CaZ06] H. Cao, X. Zhu, *A complete proof of the Poincaré and geometrization conjectures— application of the Hamilton-Perelman theory of the Ricci flow* Asian J. Math. **10** (2006), no. 2, 165–492.
- [CCR01] J. Cao, J. Cheeger and X. Rong, *Splittings and Cr-structures for manifolds with nonpositive sectional curvature*. Invent. Math. **144** (2001), no. 1, 139–167.
- [CCR04] J. Cao, J. Cheeger and X. Rong, *Local splitting structures on nonpositively curved manifolds and semirigidity in dimension 3*. Comm. Anal. Geom. **12** (2004), no. 1-2, 389–415.
- [CDM09] J. Cao, Bo Dai and Jiaqiang Mei, *An optimal extension of Perelman's comparison theorem for quadrangles and its applications*. Advanced Lectures in Mathematics, volume 11 (2009), page 39-59, In book “Recent Advances in Geometric Analysis”, edited by Y. Lee, C-S Lin, M-P Tsui. ISBN: 978-7-04-027602-2, 229 pages, Higher Educational Press and International Press.
- [CG72] J. Cheeger, D. Gromoll, *On the structure of complete manifolds of nonnegative curvature*, Annals of Math., vol **96**, (1972), 413–443.
- [CG86] J. Cheeger, M. Gromov, *Collapsing Riemannian manifolds while keeping their curvature bounded. I*. J. Differential Geom. **23** (1986), no. 3, 309–346.
- [CG90] J. Cheeger, M. Gromov, *Collapsing Riemannian manifolds while keeping their curvature bounded. II*. J. Differential Geom. **32**(1990), 269-298.
- [CM05] T. Colding, W. Minicozzi II, *Estimates for the extinction time for the Ricci flow on certain 3-manifolds and a question of Perelman*, J. Amer. Math. Soc. **18** (2005), no. 3, 561–569.

- [CM08a] T. Colding, W. Minicozzi II, *Width and mean curvature flow*, Geom. Topol. **12** (2008), no. 5, 2517–2535.
- [CM08b] T. Colding, W. Minicozzi II, *Width and finite extinction time of Ricci flow*, Geom. Topol. **12** (2008), no. 5, 2537–2586.
- [Gro81] M. Gromov, *Curvature, diameter and Betti numbers*, Comment. Math. Helv., vol **56** (1981), no. 2, 179–195.
- [Grv93] K. Grove, *Critical point theory for distance functions. Differential geometry*, In book “Riemannian geometry” (Los Angeles, CA, 1990), 357–385, Proc. Sympos. Pure Math., vol 54, Part 3, Amer. Math. Soc., Providence, RI, 1993.
- [GS77] K. Grove and K. Shiohama, *A generalized sphere theorem*, Ann of Math, **106** (1977), 201–211.
- [GW97] K. Grove, F. Wilhelm, *Metric constraints on exotic spheres via Alexandrov geometry*, J. Reine Angew. Math. **487** (1997), 201–217.
- [Ham86] R. S. Hamilton, *Four-manifolds with positive curvature operator*. J. Differential Geom. **24** (1986), no. 2, 153–179.
- [Ham99] R. S. Hamilton, *Non-singular solutions of the Ricci flow on three-manifolds*. Comm. Anal. Geom. **7** (1999), no. 4, 695–729.
- [Kap02] V. Kapovitch, *Regularity of limits of noncollapsing sequences of manifolds*, Geom. Funct. Anal. **12** (2002), no. 1, 121–137.
- [Kap05] V. Kapovitch, *Restrictions on collapsing with a lower sectional curvature bound*, Math. Z. **249**(2005), no. 3, 519–539
- [Kap07] V. Kapovitch, *Perelman’s Stability Theorem*, in “Surveys in differential geometry”, Surveys in Differential Geometry, volume **11**. International Press, Somerville, MA, 2007. xii+347 pp. ISBN: 978-1-57146-117-9, pages 103–136.
- [KPT09] V. Kapovitch, A. Petrunin, W. Tuschmann, *Nilpotency, Almost Nonnegative Curvature and Gradient Flow on Alexandrov spaces*, arXiv:math/0506273v4 [math.DG], to be appear in Annals of Mathematics.
- [KL08] B. Kleiner, J. Lott, *Notes on Perelman’s papers*, Geometry and Topology, **12** (2008) 2587–2855.
- [KL10] B. Kleiner, J. Lott, *Locally Collapsed 3-Manifolds*, in preparation.
- [MT07] J. Morgan, G. Tian, *Ricci Flow and the Poincaré Conjecture*, Clay Mathematics Monographs, vol.**3**, American Math. Society, Providence, 521 pages, 2007.
- [MT08] J. Morgan, G. Tian, *Completion of the Proof of the Geometrization Conjecture*, arXiv:0809.4040v1 [math.DG]
- [MT10] J. Morgan, G. Tian, *Ricci flow and the Geometrization Conjecture*, in preparation.
- [Per91] G. Perelman, *A. D. Alexandrov’s spaces with curvatures bounded from below. II*, Preprint.
- [Per94] G. Perelman, *Elements of Morse theory on Aleksandrov spaces*, (Russian. Russian summary) Algebra i Analiz **5** (1993), no. 1, 232–241; translation in St. Petersburg Math. J. **5** (1994), no. 1, 205–213.
- [Per97] G. Perelman, *Collapsing with no proper extremal subsets*, Comparison geometry (Berkeley, CA, 1993–94), 149–155, Math. Sci. Res. Inst. Publ., **30**, Cambridge Univ. Press, Cambridge, 1997.
- [Per02] G. Perelman, *The entropy formula for the Ricci flow and its geometric applications*, arXiv:math/0211159v1 [math.DG].

- [Per03a] G. Perelman, *Ricci flow with surgery on three-manifolds*, arXiv:math/0303109v1 [math.DG].
- [Per03b] G. Perelman, *Finite extinction time for the solutions to the Ricci flow on certain three-manifolds*, arXiv:math/0307245v1 [math.DG].
- [PP94] G. Perelman, A. Petrunin, *Extremal subsets in Aleksandrov spaces and the generalized Liberman theorem*, *Algebra i Analiz* **5** (1993), no. 1, 242–256; translation in *St. Petersburg Math. J.* **5** (1994), no. 1, 215–227.
- [Petr07] A. Petrunin, *Semiconcave functions in Alexandrov's geometry*. Surveys in differential geometry. Surveys in Differential Geometry, volume **11**. International Press, Somerville, MA, 2007. xii+347 pp. ISBN: 978-1-57146-117-9, pages 137–201.
- [Rong93] X. Rong, *The limiting eta invariants of collapsed three-manifolds*. *J. Differential Geom.* **37** (1993), no. 3, 535–568.
- [Shen96] Z. Shen, *Complete manifolds with nonnegative Ricci curvature and large volume growth*, *Invent. Math.* **125** (1996), no. 3, 393–404.
- [Shio94] T. Shioya, *Mass of rays in Alexandrov spaces of nonnegative curvature*, *Comment. Math. Helv.* **69** (1994), no. 2, 208–228.
- [SY00] T. Shioya, T. Yamaguchi, *Collapsing three-manifolds under a lower curvature bound*, *J. Differential Geom.* **56**(2000),1-66.
- [SY05] T. Shioya, T. Yamaguchi, *Volume collapsed three-manifolds with a lower curvature bound*, *Math. Ann.* **333**(2005), 131-155.
- [Wu79] H. Wu, *An elementary method in the study of non-negative curvature*, *Acta Math.* vol. **142** (1979), 57-78.
- [Yama09] T. Yamaguchi, *Collapsing and essential covering*, preprint 2009.

Jianguo Cao and Jian Ge
 Department of Mathematics
 University of Notre Dame
 Notre Dame, IN 46556, USA
 jcao@nd.edu and jge@nd.edu